arXiv:1507.06441v1 [math.SP] 23 Jul 2015

SCATTERING ON PERIODIC METRIC GRAPHS

EVGENY KOROTYAEV AND NATALIA SABUROVA

Abstract. We consider Schrödinger operators with real integrable potentials on periodic metric graphs. We show that the wave operators exist and are complete. Furthermore, we prove that the standard Fredholm determinant is well-defined without any modification for any dimension. Moreover, the determinant is analytic in the upper half-plane and the corresponding S-matrix satisfies the Birman-Krein identity. The proof is based on a fiber direct integral for metric Laplacians and a detailed analysis of the eigenvalues and the corresponding eigenfunctions of the fiber operators. In particular, we show that all eigenfunctions of the Laplacian are uniformly bounded.

Table of Contents

1. Introduction and main results 1
2. Floquet decomposition of metric Laplacians 10
3. Eigenfunctions of metric Laplacians 13
4. Schrödinger operators on periodic metric graphs 25
5. Lattice graph 29
6. Graphene lattice 32
7. Stanene lattice 34

1. Introduction and main results

1.1. Introduction. We consider a Schrödinger operator $H = H_0 + Q$ on $\mathbb{Z}^d$-periodic metric graphs Γ with unit-length edges, i.e., on the so-called $\mathbb{Z}^d$-periodic equilateral graphs, $d \geqslant 2$. Here $H_0 = \Delta_M$ is a positive metric Laplace operator with Kirchhoff vertex conditions and Q is the real integrable potential. Differential operators on metric graphs arise naturally as simplified models in mathematics, physics, chemistry, and engineering.

It is well-known that Δ_M is self-adjoint and its spectrum consists of an absolutely continuous part plus an infinite number of flat bands (i.e., eigenvalues with infinite multiplicity). The absolutely continuous spectrum is a union of an infinite number of spectral bands separated by gaps. The spectrum of metric Laplacians has deep relationships with the spectrum of discrete Laplacians Δ on the corresponding discrete graphs, see [B85], [BGP08], [C97] and references therein.

There are a lot of papers, and even books, on the spectrum of discrete and metric Laplacians on finite and infinite graphs (see [BK13], [Ch97], [CDS95], [CDGT88], [P12] and references therein).

There are results about spectral properties of discrete Schrödinger operators on the lattice $\mathbb{Z}^d$, the simplest example of $\mathbb{Z}^d$-periodic graphs. Schrödinger operators with decaying

---

Date: July 24, 2015.

Key words and phrases. direct integral, scattering, Fredholm determinant, metric Laplacian, Schrödinger operators, periodic metric graph.

potentials on the lattice $\mathbb{Z}^d$ are considered by Boutet de Monvel-Sahbani [BS99], Hundertmark-Simon [HS02], Isozaki-Korotyaev [IK12], Isozaki-Morioka [IM14], Nakamura [Na14], Rosenblum-Solomjak [RoS09], Shaban-Vainberg [SV01] and see references therein. Gieseker-Knörrer-Trubowitz [GKT93] consider Schrödinger operators with periodic potentials on the lattice $\mathbb{Z}^2$. Korotyaev-Kutsenko [KK10] study the spectra of the discrete Schrödinger operators on graphene nano-ribbons in external electric fields. Ando [A12] considers the inverse spectral theory for the discrete Schrödinger operators with finitely supported potentials on the hexagonal lattice.

The decomposition of discrete Schrödinger operators with periodic potentials on periodic graphs into a constant fiber direct integral (1.4) (without an exact form of fiber operators) is discussed by Higuchi-Shirai [HS04], Higuchi-Nomura [HN09], Rabinovich-Roch [RR07]. In particular, they prove that the spectrum of Schrödinger operators consists of an absolutely continuous part and a finite number of flat bands (i.e., eigenvalues with infinite multiplicity). The absolutely continuous spectrum consists of a finite number of bands (intervals) separated by gaps. The fiber operator is determined by Korotyaev-Saburova [KS14a]. Moreover, they estimate the Lebesgue measure of the spectrum of the discrete Schrödinger operator in terms of geometric parameters of the graph and the potential. Furthermore, they estimate a global variation of the Lebesgue measure of the spectrum and a global variation of gap-length in terms of potentials and geometric parameters of the graph.

Lledó-Post [LP08], Korotyaev-Saburova [KS14b], [KS15a] describe a localization of spectral bands of Laplacians both on metric and discrete periodic graphs in terms of eigenvalues of the operator on finite graphs (the so-called eigenvalue bracketing).

Korotyaev-Saburova [KS15b] consider Laplacians on periodic equilateral metric graphs and estimate the Lebesgue measure of the bands and gaps on a finite interval in terms of geometric parameters of the graph.

Effective masses for Laplacians on periodic discrete and metric equilateral graphs are studied by Korotyaev-Saburova [KS15c]. They estimate effective masses associated with the ends of each spectral band in terms of geometric parameters of graphs. Moreover, at the beginning of the spectrum they obtain two-sided estimates of the effective mass in terms of geometric parameters of graphs. These results are essentially used in this paper.

We do not know any papers about scattering on periodic metric graphs. Note that there are some results about the scattering on a graph that consists of a compact part with one or more infinite leads attached to it (see the survey in Section 5.5 in [BK13]).

We describe our main results:

i) The direct integral for a metric Laplacian Δ_M *is described in terms of the discrete Laplacian* Δ.

ii) Eigenfunctions of fiber operators for the metric Laplacian with eigenvalues from the set $\sigma(\Delta_M) \setminus \sigma_D$ *are described in terms of eigenfunctions of fiber operators for the discrete Laplacian* Δ*, where* $\sigma_D = \{(\pi j)^2 : j \in \mathbb{N}\}$.

iii) Eigenfunctions of fiber operators for the metric Laplacian corresponding to the flat bands $(\pi j)^2$, $j \in \mathbb{N}$, *(the so-called Dirichlet spectrum) are determined.*

iv) All eigenfunctions of the metric Laplacian are uniformly bounded;

v) We consider scattering for a Schrödinger operator $H = \Delta_M + Q$*, where the potential* $Q \in L^1(\Gamma)$ *is real. In particular, we obtain*

- *the existence and completeness of the wave operators,*

- *a standard Fredholm determinant (without any modification for any dimension) is well defined and is analytic in the upper half-plane and its main properties are discussed.*
- *the difference of the resolvents for a Schrödinger operator and for the Laplace operator belongs to the trace class for any dimension. Note that for corresponding Schrödinger operators on $\mathbb{R}^d$, $d \geqslant 2$, it does not hold true.*

The proof of these results is based on a detailed analysis of eigenfunctions of fiber operators for the metric Laplacian obtained in the present paper. Note that for short range potentials one needs to develop the Mourre's or Enss's approaches on periodic metric graphs.

1.2. **Metric Laplacians.** Let $\Gamma = (V, \mathcal{E})$ be a connected infinite graph, possibly having loops and multiple edges, where V is the set of its vertices and $\mathcal{E}$ is the set of its unoriented edges. An edge connecting vertices u and v from V will be denoted as the unordered pair $(u, v)_e \in \mathcal{E}$ and is said to be *incident* to the vertices. Vertices $u, v \in V$ will be called *adjacent* and denoted by $u \sim v$, if $(u, v)_e \in \mathcal{E}$. Denote by $I(v)$ the set of all edges of Γ incident to the vertex $v \in V$. We define the degree $\varkappa_v$ of the vertex $v \in V$ as the number of all its incident edges from $\mathcal{E}$ (here a loop is counted twice).

Below we consider locally finite $\mathbb{Z}^d$-periodic metric equilateral graphs Γ, i.e., graphs satisfying the following conditions:

1) Γ is equipped with an action of the free abelian group $\mathbb{Z}^d$;
2) the degree of each vertex is finite;
3) the quotient graph $\Gamma_* = \Gamma/\mathbb{Z}^d$ is compact;
4) all edges of the graph are assigned unit length.

Remark. 1) We do not assume the graph be embedded into an Euclidean space. But in some application such a natural embedding exists (e.g., in modeling waves in thin branching "graph-like" structures: narrow waveguides, quantum wires, photonic crystal, blood vessels, lungs, see [BK13], [P12]). In this case a simple geometric model is a graph Γ embedded into $\mathbb{R}^d$ in such a way that it is invariant with respect to the shifts by integer vectors $m \in \mathbb{Z}^d$, which produce an action of $\mathbb{Z}^d$.

2) We also call the quotient graph $\Gamma_* = \Gamma/\mathbb{Z}^d$ *the fundamental graph* of the periodic graph Γ. If Γ is embedded into the space $\mathbb{R}^d$ the fundamental graph Γ_* is a graph on the surface $\mathbb{R}^d/\mathbb{Z}^d$. The fundamental graph $\Gamma_* = (V_*, \mathcal{E}_*)$ has the vertex set $V_* = V/\mathbb{Z}^d$ and the set $\mathcal{E}_* = \mathcal{E}/\mathbb{Z}^d$ of unoriented edges. Let $\nu = \# V_*$, where $\# A$ is the number of elements of the set A. Denote by $I_*(v)$ the set of all edges of Γ_* incident to the vertex $v \in V_*$.

Each edge $\mathbf{e}$ of Γ will be identified with the segment $[0, 1]$. This identification introduces a local coordinate $t \in [0, 1]$ along each edge. Thus, we give an orientation on the edge. An edge starting at $u \in V$ and ending at $v \in V$ will be denoted as the ordered pair $(u, v) \in \mathcal{E}$. Note that the spectrum of Laplacians on metric graphs does not depend on the orientation of graph edges. For each function y on Γ we define a function $y_{\mathbf{e}} = y\big|_{\mathbf{e}}$, $\mathbf{e} \in \mathcal{E}$. We identify each function $y_{\mathbf{e}}$ on $\mathbf{e}$ with a function on $[0, 1]$ by using the local coordinate $t \in [0, 1]$. Let $L^2(\Gamma) = \oplus_{\mathbf{e} \in \mathcal{E}} L^2(\mathbf{e})$ be the Hilbert space of all functions $y = (y_{\mathbf{e}})_{\mathbf{e} \in \mathcal{E}}$, where each $y_{\mathbf{e}} \in L^2(\mathbf{e}) = L^2(0, 1)$, equipped with the norm

$$\|y\|^2_{L^2(\Gamma)} = \sum_{\mathbf{e} \in \mathcal{E}} \|y_{\mathbf{e}}\|^2_{L^2(0,1)} < \infty.$$

We define the positive metric Laplacian Δ_M on $y = (y_{\mathbf{e}})_{\mathbf{e} \in \mathcal{E}} \in L^2(\Gamma)$ by

$$(\Delta_M y)_{\mathbf{e}} = -y''_{\mathbf{e}}, \qquad \text{where} \qquad (y''_{\mathbf{e}})_{\mathbf{e} \in \mathcal{E}} \in L^2(\Gamma),$$

where y satisfies the Kirchhoff conditions:

$$y \text{ is continuous on } \Gamma, \qquad \sum_{\mathbf{e}\in I(v)} (-1)^{\delta(\mathbf{e},v)}\, y'_{\mathbf{e}}\big(\delta(\mathbf{e},v)\big)=0, \quad \forall v\in V, \tag{1.1}$$

$$\delta(\mathbf{e},v)=\left\{\begin{array}{ll} 1, & \text{if } v \text{ is the terminal vertex of the edge } \mathbf{e}\\ 0, & \text{if } v \text{ is the initial vertex of the edge } \mathbf{e}.\end{array}\right. \tag{1.2}$$

1.3. **Discrete Laplacians.** In order to define the Floquet-Bloch decomposition (1.4) of metric Laplacians we need to introduce the inverse edge (v,u) for each oriented edge $(u,v)\in\mathcal{E}$. Let $\mathcal{A}$ and $\mathcal{A}_*$ denote the sets of all oriented edges and their inverse edges of the periodic graph Γ and its fundamental graph Γ_*, respectively.

Let $\ell^2(V)$ be the Hilbert space of all functions $f:V\to\mathbb{C}$, equipped with the norm

$$\|f\|^2_{\ell^2(V)}=\sum_{v\in V}|f(v)|^2<\infty.$$

We define the discrete normalized Laplacian (i.e., the Laplace operator) Δ on $f\in\ell^2(V)$ by

$$\big(\Delta f\big)(v)=-\frac{1}{\sqrt{\varkappa_v}}\sum_{(v,\,u)_e\in\mathcal{E}}\frac{1}{\sqrt{\varkappa_u}}\,f(u), \qquad v\in V, \tag{1.3}$$

where $\varkappa_v$ is the degree of the vertex $v\in V$ and all loops in the sum (1.3) are counted twice. This Laplacian Δ on $\ell^2(V)$ is self-adjoint and has the standard decomposition into a constant fiber direct integral by

$$\ell^2(V)=\frac{1}{(2\pi)^d}\int^{\oplus}_{\mathbb{T}^d}\ell^2(V_*)\,d\vartheta, \qquad U\Delta U^{-1}=\frac{1}{(2\pi)^d}\int^{\oplus}_{\mathbb{T}^d}\Delta(\vartheta)d\vartheta, \tag{1.4}$$

$\mathbb{T}^d=\mathbb{R}^d/(2\pi\mathbb{Z})^d$, for some unitary operator U. Here $\ell^2(V_*)=\mathbb{C}^\nu$ is the fiber space and $\Delta(\vartheta)$ is the $\nu\times\nu$ Floquet (fiber) matrix given by (2.5), where $\nu=\#V_*$. The Floquet matrix $\Delta(\vartheta)$ has ν eigenvalues. Recall that λ_* is an eigenvalue of Δ iff λ_* is an eigenvalue of $\Delta(\vartheta)$ for any $\vartheta\in\mathbb{T}^d$ (see Proposition 4.2 in [HN09]). Thus, we can define the multiplicity of a flat band by: an eigenvalue λ_* of Δ has the multiplicity m iff $\lambda_*=\text{const}$ is an eigenvalue of $\Delta(\vartheta)$ with the multiplicity m for almost all $\vartheta\in\mathbb{T}^d$. Thus, if the operator Δ has $r\geqslant 0$ flat bands, then we denote them by

$$\lambda_j(\vartheta)=\text{const}, \qquad \forall\,\nu-r<j\leqslant\nu=\#V_*, \quad \forall\,\vartheta\in\mathbb{T}^d. \tag{1.5}$$

Here we separate all flat bands from other bands of the Laplacian. All other eigenvalues $\lambda_1(\vartheta),\ldots,\lambda_{\nu-r}(\vartheta)$ are not constant. They can be enumerated in increasing order (counting multiplicities) by

$$\lambda_1(\vartheta)\leqslant\lambda_2(\vartheta)\leqslant\ldots\leqslant\lambda_{\nu-r}(\vartheta), \qquad \forall\,\vartheta\in\mathbb{T}^d. \tag{1.6}$$

Since $\Delta(\vartheta)$ is self-adjoint and analytic in $\vartheta\in\mathbb{T}^d$, each $\lambda_n(\cdot)$, $n\in\mathbb{N}_\nu=\{1,\ldots,\nu\}$, is a real and piecewise analytic function on the torus $\mathbb{T}^d$. We denote the corresponding projector by $P_n(\vartheta)$ and we have

$$\Delta(\vartheta)=\sum_{n=1}^{\nu}\lambda_n(\vartheta)P_n(\vartheta), \qquad \vartheta\in\mathbb{T}^d. \tag{1.7}$$

We define the spectral bands $\sigma_n(\Delta)$ by

$$\sigma_n(\Delta)=[\lambda_n^-,\lambda_n^+]=\lambda_n(\mathbb{T}^d), \qquad n\in\mathbb{N}_\nu. \tag{1.8}$$

Thus, the spectrum of the Laplacian Δ on Γ has the form $\sigma(\Delta)=\bigcup_{n=1}^{\nu}\sigma_n(\Delta)$ and we get

$$\sigma(\Delta)=\sigma_{ac}(\Delta)\cup\sigma_{fb}(\Delta),\qquad \sigma_{ac}(\Delta)=\bigcup_{n=1}^{\nu-r}\sigma_n(\Delta),\qquad \sigma_{fb}(\Delta)=\bigcup_{n=\nu-r+1}^{\nu}\sigma_n(\Delta). \tag{1.9}$$

Here $\sigma_{ac}(\Delta)$ is the absolutely continuous spectrum, which is a union of non-degenerate intervals from (1.8), and $\sigma_{fb}(\Delta)$ is the set of all flat bands (eigenvalues of infinite multiplicity). An open interval between two neighboring non-degenerate spectral bands is called a *spectral gap*.

1.4. **Direct integral of metric Laplacians.** We introduce the Hilbert space (a constant fiber direct integral)

$$\mathscr{H}=L^2\big(\mathbb{T}^d,\frac{d\vartheta}{(2\pi)^d},\mathcal{H}\big)=\int_{\mathbb{T}^d}^{\oplus}\mathcal{H}\,\frac{d\vartheta}{(2\pi)^d}\,,\qquad \mathcal{H}=L^2(\Gamma_*),$$

equipped with the norm

$$\|g\|^2_{\mathscr{H}}=\int_{\mathbb{T}^d}\|g(\vartheta,\cdot)\|^2_{\mathcal{H}}\frac{d\vartheta}{(2\pi)^d}\,,$$

where a function $g(\vartheta,\cdot)\in L^2(\Gamma_*)$ for all $\vartheta\in\mathbb{T}^d$. We have the preliminary standard result about a direct integral, see, e.g., [RS78].

Theorem 1.1. *The metric Laplacian Δ_M on $L^2(\Gamma)$ has the following decomposition into a constant fiber direct integral*

$$L^2(\Gamma)=\frac{1}{(2\pi)^d}\int_{\mathbb{T}^d}^{\oplus}L^2(\Gamma_*)\,d\vartheta,\qquad \mathscr{U}\Delta_M\mathscr{U}^{-1}=\frac{1}{(2\pi)^d}\int_{\mathbb{T}^d}^{\oplus}\Delta_M(\vartheta)d\vartheta, \tag{1.10}$$

for the unitary operator $\mathscr{U}:L^2(\Gamma)\to\mathscr{H}$ defined by

$$(\mathscr{U}h)(\vartheta,x)=\sum_{m\in\mathbb{Z}^d}e^{-i\langle m,\vartheta\rangle}h(x+m),\qquad (\vartheta,x)\in\mathbb{T}^d\times\Gamma_*, \tag{1.11}$$

where $x+m$ denotes the action of $m\in\mathbb{Z}^d$ on $x\in\Gamma_*$. Here the Floquet (fiber) operator $\Delta_M(\vartheta)$ acts on $y=(y_{\mathbf{e}})_{\mathbf{e}\in\mathcal{E}_*}\in L^2(\Gamma_*)$ by

$$(\Delta_M(\vartheta)y)_{\mathbf{e}}=-y''_{\mathbf{e}},\qquad (y''_{\mathbf{e}})_{\mathbf{e}\in\mathcal{E}_*}\in L^2(\Gamma_*), \tag{1.12}$$

and the function y satisfies the so-called quasi-periodic conditions at any $v\in V_*$:

$$e^{-i\delta(\mathbf{e}_1,v)\langle\tau(\mathbf{e}_1),\,\vartheta\rangle}y_{\mathbf{e}_1}\big(\delta(\mathbf{e}_1,v)\big)=e^{-i\delta(\mathbf{e}_2,v)\langle\tau(\mathbf{e}_2),\,\vartheta\rangle}y_{\mathbf{e}_2}\big(\delta(\mathbf{e}_2,v)\big),\quad \forall\,\mathbf{e}_1,\mathbf{e}_2\in I_*(v), \tag{1.13}$$

$$\sum_{\mathbf{e}\in I_*(v)}(-1)^{\delta(\mathbf{e},v)}e^{-i\delta(\mathbf{e},v)\langle\tau(\mathbf{e}),\,\vartheta\rangle}\,y'_{\mathbf{e}}\big(\delta(\mathbf{e},v)\big)=0, \tag{1.14}$$

where $\delta(\mathbf{e},v)$ is defined by (1.2), $\tau(\mathbf{e})\in\mathbb{Z}^d$ is the edge index, defined in subsection 2.1, and $\langle\cdot\,,\cdot\rangle$ denotes the standard inner product in $\mathbb{R}^d$.

Remark. In (1.11) we identify an edge $\mathbf{e}_*\in\mathcal{E}_*$ of the fundamental graph Γ_* with an edge $\mathbf{e}\in\mathcal{E}$ of the graph Γ (for more details see Subsection 2.2).

1.5. **Spectrum of the metric Laplacian Δ_M and its fiber operators $\Delta_M(\vartheta)$.** Consider the eigenvalues problem on the unit interval with Dirichlet boundary conditions

$$-y''=Ey,\qquad y(0)=y(1)=0. \tag{1.15}$$

It is known that each $(\pi j)^2$, $j\in\mathbb{N}$, is the so-called Dirichlet eigenvalue of the problem (1.15). We define its spectrum by

$$\sigma_D=\{(\pi j)^2 : j\in\mathbb{N}\}. \tag{1.16}$$

We describe our first main results about eigenfunctions of the metric Laplacian Δ_M having eigenvalues $(\pi j)^2, j\in\mathbb{N}$.

Theorem 1.2. *Let $\Delta_M(\vartheta)$, $\vartheta\in\mathbb{T}^d\setminus\{0\}$, be the Floquet operator on the fundamental graph Γ_*. Then for all $j\in\mathbb{N}$ the following statements hold true:*

i) $E_j^0=(\pi j)^2$ is the eigenvalue of the operator $\Delta_M(\vartheta)$.

ii) Each eigenvalue $(\pi j)^2$ has multiplicity $\nu_*-\nu\geqslant d-1$, where $\nu=\#V_*$ is the number of the fundamental graph vertices and $\nu_*=\#\mathcal{E}_*$ is the number of the fundamental graph edges from $\mathcal{E}_*$.

iii) Corresponding normalized eigenfunctions $\Psi_{s,j}^0=\big(\Psi_{s,j,\mathbf{e}}^0\big)_{\mathbf{e}\in\mathcal{E}_*}$, $s\in\mathbb{N}_{\nu_*-\nu}$, have the form:

$$\begin{aligned}\Psi_{s,j,\mathbf{e}}^0(\vartheta,t)=X_{s,j,\mathbf{e}}(\vartheta)\sqrt{2}\,\sin(\pi jt),\quad t\in[0,1],\\ \|\Psi_{s,j}^0(\vartheta,\cdot)\|^2_{L^2(\Gamma_*)}=\sum_{\mathbf{e}\in\mathcal{E}_*}|X_{s,j,\mathbf{e}}(\vartheta)|^2=1,\end{aligned} \tag{1.17}$$

where $\big(X_{s,j,\mathbf{e}}(\vartheta)\big)_{\mathbf{e}\in\mathcal{E}_*}$ is a normalized solution of the system of ν equations

$$\sum_{\substack{\mathbf{e}\in\mathcal{E}_*,\\ \text{starting at } v}} x_{\mathbf{e}}-(-1)^j\sum_{\substack{\mathbf{e}\in\mathcal{E}_*,\\ \text{ending at } v}} e^{-i\langle\tau(\mathbf{e}),\,\vartheta\rangle}\,x_{\mathbf{e}}=0,\qquad \forall v\in V_*, \tag{1.18}$$

one equation for each $v\in V_*$. This system has the rank ν and thus there exist $\nu_*-\nu$ linearly independent normalized solutions $\big(X_{s,j,\mathbf{e}}(\vartheta)\big)_{\mathbf{e}\in\mathcal{E}_*}$, $s\in\mathbb{N}_{\nu_*-\nu}$. Moreover, $X_{s,j,\mathbf{e}}(\vartheta)=X_{s,j+2,\mathbf{e}}(\vartheta)$ for all $(s,\mathbf{e})\in\mathbb{N}_{\nu_*-\nu}\times\mathcal{E}_*$ and the eigenfunctions $\Psi_{s,j}^0(\vartheta,t)$ satisfy

$$\sup_{(s,j)\in\mathbb{N}_{\nu_*-\nu}\times\mathbb{N}}\|\Psi_{s,j}^0(\vartheta,\cdot)\|_{L^\infty(\Gamma_*)}\leqslant\sqrt{2}\,. \tag{1.19}$$

Remark. 1) In order to determine $X_{s,j,\mathbf{e}}$ in (1.17) we have the $\nu\times\nu_*$ linear system (1.18) with ν_* variables.

2) The eigenfunctions $\Psi_{s,j}^0$ defined by (1.17) vanish at each vertex of the fundamental graph.

3) The Floquet operator $\Delta_M(0)$ is exactly the metric Laplacian on the fundamental graph Γ_*. Moreover, it is known (see e.g., [B85], [C97], [LP08]) that

i) $(\pi j)^2$, $j\in\mathbb{N}$, is the eigenvalue of the operator $\Delta_M(0)$;

ii) each eigenvalue $(2\pi j)^2$ has multiplicity $\nu_*-\nu+2$;

iii) each eigenvalue $(2j+1)^2\pi^2$ has multiplicity $\nu_*-\nu+2$, if Γ_* is bipartite and multiplicity $\nu_*-\nu$, if Γ_* is not bipartite;

iv) the corresponding eigenfunctions are also constructed in the papers mentioned above (for more details see Remark after the proof of Theorem 1.2).

4) The number of vertices ν and the number of edges ν_* of the fundamental graph can be arbitrary, but there is a restriction (see Theorem 1.2 in [KS15a]):

$$\nu+d\leqslant\nu_*+1.$$

Theorem 1.2 allows to write the operator Δ_M in the form $\Delta_M = \Delta_{MD} \oplus \Delta_{MV}$, where

$$\Delta_M(\vartheta) = \Delta_{MD}(\vartheta) \oplus \Delta_{MV}(\vartheta), \qquad \Delta_{MD}(\vartheta) = \sum_{j\geqslant 1} (\pi j)^2 \mathcal{P}_j^0(\vartheta), \tag{1.20}$$

$\mathcal{P}_j^0(\vartheta)$ is the projector corresponding to the Dirichlet eigenvalue $(\pi j)^2$ and its eigenfunctions $\Psi_{s,j}^0$, where $(s,j) \in \mathbb{N}_{\nu_*-\nu} \times \mathbb{N}$. The operator $\Delta_{MD}(\vartheta)$ is associated with the Dirichlet eigenvalues σ_D. The other part $\Delta_{MV}(\vartheta)$ is associated with the vertex set V_* and we discuss one in the next theorem. Below without loss of generality we assume that for a bipartite periodic graph Γ the fundamental graph Γ_* is also bipartite.

We describe our second main results about the "vertex Laplacian" Δ_{MV} and its eigenfunctions. These eigenfunctions are expressed in terms of eigenfunctions of discrete Laplacians.

Theorem 1.3. *The operator $\Delta_{MV}(\vartheta)$ on Γ_* defined by (1.20) has a spectral representation given by*

$$\Delta_{MV}(\vartheta) = \sum_{(n,j)\in\mathbb{N}_\nu\times\mathbb{Z}_0} z_{n,j}^2(\vartheta)\mathcal{P}_{n,j}(\vartheta), \qquad \textit{for all} \quad \vartheta \in \mathbb{T}^d \setminus \{0\}, \tag{1.21}$$

where its eigenvalues $z_{n,j}^2(\vartheta)$ and the eigenfunctions $\Psi_{n,j}(\vartheta) = \big(\Psi_{n,j,\mathbf{e}}(\vartheta,t)\big)_{\mathbf{e}\in\mathcal{E}_*}$ corresponding to the projectors $\mathcal{P}_{n,j}(\vartheta)$ satisfy:

$$z_{n,j}(\vartheta) = \begin{cases} z_n(\vartheta) + \pi j, & j \textit{ is even} \\ (\pi - z_n(\vartheta)) + \pi j, & j \textit{ is odd} \end{cases}, \quad z_n(\vartheta) = \arccos(-\lambda_n(\vartheta)) \in (0,\pi) \tag{1.22}$$

and (here $\mathbf{e} = (u,v)$ and $n \in \mathbb{N}_\nu$, $j \in \mathbb{Z}_0 = \{0,1,2,\ldots\}$)

$$\Psi_{n,j,\mathbf{e}}(\vartheta,t) = \tfrac{\sqrt{2}}{\sin z_n(\vartheta)} \Big(\tfrac{\psi_n(\vartheta,u)}{\sqrt{\varkappa_u}} \sin(z_{n,j}(\vartheta)\,(1-t)) + \tfrac{\psi_n(\vartheta,v)}{\sqrt{\varkappa_v}} e^{i\langle\tau(\mathbf{e}),\,\vartheta\rangle} \sin(z_{n,j}(\vartheta)\,t) \Big), \tag{1.23}$$

$$\|\Psi_{n,j}(\vartheta)\|_{L^2(\Gamma_*)} = \|\psi_n(\vartheta,\cdot)\|_{\ell^2(V_*)} = 1, \tag{1.24}$$

$$\sup_{(n,j,\vartheta)\in\mathbb{N}_\nu\times\mathbb{Z}_0\times\mathbb{T}^d} \|\Psi_{n,j}(\vartheta)\|_{L^\infty(\Gamma_*)} = C < \infty, \tag{1.25}$$

where $\psi_n(\vartheta,\cdot) \in \ell^2(V_*)$ is the normalized eigenfunction of the operator $\Delta(\vartheta)$ corresponding to the eigenvalue $\lambda_n(\vartheta)$.

Remark. 1) The eigenvalue $z_{n,j}^2(\vartheta) \neq (\pi j)^2$ for all $\vartheta \in \mathbb{T}^d \setminus \{0\}$ and all $(n,j) \in \mathbb{N}_\nu \times \mathbb{Z}_0$.

2) Recently, Pankrashkin (see Proposition 1 in [Pa12], Theorem 14 in [Pa13]) shows the following result: Let $\eta(z) = -\cos\sqrt{z}$ and χ_ω be the characteristic function of the set $\omega \subset \mathbb{R}$ and let an interval $J \subset \mathbb{R} \setminus \sigma_D$. Then operator $A = \Delta_{MV}\chi_J(\Delta_{MV})$ is unitary equivalent to the operator $B = \eta^{-1}(\Delta\chi_{\eta(J)}(\Delta))$, i.e., the following identity $A = UBU^{-1}$ holds true for some unitary operator U.

3) We do not know any paper about (1.23) – (1.25). These results are important to study the Fredholm determinant in Theorem 1.6.

4) The explicit form of the constant C in (1.25) is defined in Propositions 3.3, 3.4. Results about the effective masses for Laplacians [KS15c] are essential in the proof of (1.25).

Theorems 1.1 – 1.3 imply the well-known results, see, e.g., [BK13], [BGP08], [P12].

Corollary 1.4. *The Laplacian Δ_M on $L^2(\Gamma)$ has the spectrum given by*

$$\begin{aligned} \sigma(\Delta_M)=\sigma_{ac}(\Delta_M)\cup\sigma_{fb}(\Delta_M),\qquad \sigma_{ac}(\Delta_M)=\bigcup_{(n,j)\in\mathbb{N}_{\nu-r}\times\mathbb{N}}\sigma_{n,j-1}(\Delta_M),\\ \sigma_{fb}(\Delta_M)=\sigma_{fb}(\Delta_{MD})\cup\sigma_{fb}(\Delta_{MV}),\qquad \sigma_{fb}(\Delta_{MD})=\sigma_D,\\ \sigma_{fb}(\Delta_{MV})=\bigcup_{n=\nu-r+1}^{\nu}\bigcup_{j=0}^{\infty}\sigma_{n,j}(\Delta_M),\qquad \sigma_{n,j}(\Delta_M)=[E_{n,j}^-,E_{n,j}^+], \end{aligned} \tag{1.26}$$

$$E_{n,j}^{\pm}=\begin{cases}\big(z_n^{\pm}+\pi j\big)^2, & j \textit{ is even}\\ \big((\pi-z_n^{\mp})+\pi j\big)^2, & j \textit{ is odd}\end{cases},\qquad z_n^{\pm}=\arccos(-\lambda_n^{\pm})\in[0,\pi],\quad n\in\mathbb{N}_\nu.$$

Here $\sigma_{ac}(\Delta_M)$ is the absolutely continuous spectrum, which is a union of non-degenerate intervals, and $\sigma_{fb}(\Delta_M)$ is the set of all flat bands (eigenvalues of infinite multiplicity).

Remarks. 1) Recall that Brüning-Geyler-Pankrashkin [BGP08] show the following identities

$$\sigma_\alpha(\Delta_M)\setminus\sigma_D=\{E\in\mathbb{R}\setminus\sigma_D:-\cos\sqrt{E}\in\sigma_\alpha(\Delta)\},\qquad \alpha\in\{\text{fb, ac}\},\tag{1.27}$$

where $\sigma_D=\{(\pi j)^2: j\in\mathbb{N}\}$.

2) The identities (1.26) show that spectral properties of the discrete Laplacian Δ are important to study spectral properties of the metric Laplacian Δ_M.

Now we discuss the connection between eigenfunctions of the absolutely continuous spectrum of the discrete and metric Laplacians and eigenfunctions of their fiber operators.

Proposition 1.5. *Let $\psi_n(\vartheta,\cdot)\in\ell^2(V_*)$ be the normalized eigenfunction of the fiber operator $\Delta(\vartheta)$ corresponding to the eigenvalue $\lambda_n(\vartheta)\in\sigma_{ac}(\Delta)$ and let $\Psi_{n,j}(\vartheta,\cdot)\in L^2(\Gamma_*)$ be the normalized eigenfunction of the fiber operator $\Delta_M(\vartheta)$ corresponding to the eigenvalue $z_{n,j}^2(\vartheta)\in\sigma_{ac}(\Delta_M)$, where $\vartheta\in\mathbb{T}^d\setminus\{0\}$ and $z_{n,j}^2(\vartheta)$ is defined by (1.22). Then the following statements hold true.*

i) The eigenfunction $\widetilde{\psi}_n(\vartheta,\cdot)$ of the absolutely continuous spectrum of the discrete Laplacian Δ corresponding to $\lambda_n(\vartheta)$ has the form

$$\widetilde{\psi}_n(\vartheta,v+m)=e^{i\langle m,\vartheta\rangle}\psi_n(\vartheta,v),\qquad \forall\,(v,m)\in V_*\times\mathbb{Z}^d.\tag{1.28}$$

ii) The eigenfunction $\widetilde{\Psi}_{n,j}(\vartheta,\cdot)$ of the absolutely continuous spectrum of the metric Laplacian Δ_M corresponding to $z_{n,j}^2(\vartheta)$ has the form

$$\widetilde{\Psi}_{n,j}(\vartheta,x+m)=e^{i\langle m,\vartheta\rangle}\Psi_{n,j}(\vartheta,x),\qquad \forall\,(x,m)\in\Gamma_*\times\mathbb{Z}^d.\tag{1.29}$$

iii) If a $\mathbb{Z}^d$-periodic graph Γ is embedded into $\mathbb{R}^d$, then the eigenfunctions from (1.28), (1.29) satisfy

$$\widetilde{\psi}_n(\vartheta,v)=e^{i\langle v,\vartheta\rangle}\varphi_n(\vartheta,v),\qquad \forall\,v\in V;\tag{1.30}$$

$$\widetilde{\Psi}_n(\vartheta,x)=e^{i\langle x,\vartheta\rangle}\Phi_n(\vartheta,x),\qquad \forall\,x\in\Gamma;\tag{1.31}$$

where the functions $\varphi_n(\vartheta,v)$ and $\Phi_n(\vartheta,x)$ are $\mathbb{Z}^d$-periodic with respect to v and x, respectively.

Remarks. 1) In (1.28), (1.29) we identify the vertices and edges of the fundamental graph Γ_* with some vertices and edges of the periodic graph Γ. For more details see the proof of this proposition.

2) From (1.29) and (1.25) it follows that all eigenfunctions of the absolutely continuous spectrum of the Laplacian on a periodic metric graph are uniformly bounded in $L^\infty(\Gamma)$.

3) Formulas (1.23), (1.28) and (1.29) give that the eigenfunctions of the absolutely continuous spectrum of the discrete and metric Laplacians are connected by the same relation (1.23) as the eigenfunctions of their fiber operators.

4) Eigenfunctions of the discrete and metric Laplacians corresponding to flat bands are connected with eigenfunctions of their fiber operators by the Floquet transforms (for more details see Proposition 3.5).

1.6. **Scattering on metric graphs.** We consider a Schrödinger operator $H = H_0 + Q$ on $L^2(\Gamma)$, where $H_0 = \Delta_M$ and the potential $Q \in L^2(\Gamma) \cap L^1(\Gamma)$ is real. Here $L^1(\Gamma)$ is the space of all functions $f = (f_{\mathbf{e}})_{\mathbf{e}\in\mathcal{E}}$ on Γ equipped with the norm $\|f\|_{L^1(\Gamma)} = \sum_{\mathbf{e}\in\mathcal{E}} \|f_{\mathbf{e}}\|_{L^1(\mathbf{e})}$. Let $\mathbf{B}_1$ and $\mathbf{B}_2$ be the trace and the Hilbert-Schmidt class equipped with the norm $\|\cdot\|_{\mathbf{B}_1}$ and $\|\cdot\|_{\mathbf{B}_2}$ correspondingly.

In Theorem 1.6 we show that $QR_0(k) \in \mathbf{B}_2$ for all $k \in \mathbb{C}_+$. This gives that the operator H is self-adjoint on $\mathscr{D}(H_0)$. For any $k \in \mathbb{C}_+$ we put

$$\begin{aligned} &R_0(k) = (H_0 - k^2)^{-1}, \qquad R(k) = (H - k^2)^{-1}, \\ &Y_0(k) = |Q|^{\frac12} R_0(k)\, Q^{\frac12}, \qquad Q^{\frac12} = |Q|^{\frac12} \operatorname{sign} Q. \end{aligned} \tag{1.32}$$

Below we show that the operator $Y_0(k)$ belongs to the trace class. Thus, we can define the Fredholm determinant D by

$$D(k) = \det(I + Y_0(k)), \qquad k \in \mathbb{C}_+. \tag{1.33}$$

We describe our third main result about scattering on metric graphs.

Theorem 1.6. *Let $Q \in L^2(\Gamma)$ be real. Then $QR_0(k) \in \mathbf{B}_2$ for all $k \in \mathbb{C}_+$ and the operator $H = H_0 + Q$ is self-adjoint on $\mathscr{D}(H_0)$. Let, in addition, $Q \in L^1(\Gamma)$. Then*

$$R(k) - R_0(k), \quad Y_0(k) \in \mathbf{B}_1 \qquad \forall\ k \in \mathbb{C}_+, \tag{1.34}$$

and the determinant $D(k) = \det(I + Y_0(k))$ is well-defined and is analytic in $\mathbb{C}_+$ and the limit $D(k + i0)$ exists for almost all $k \in \mathbb{R}$. Furthermore, the wave operators

$$W_\pm = \text{s-}\lim e^{itH} e^{-itH_0} P_{ac}(H_0) \qquad as \qquad t \to \pm\infty, \tag{1.35}$$

exist and are complete, i.e. the range of $W_\pm$ is equal to $\mathscr{H}_{ac}(H)$ and $\sigma_{ac}(H) = \sigma_{ac}(H_0)$. Moreover, the S-operator given by

$$S = W_+^* W_- \tag{1.36}$$

is unitary on $\mathscr{H}_{ac}(H_0)$ and the corresponding S-matrix $S(k)$ (defined by (1.39)) for almost all $k^2 \in \sigma_{ac}(H_0)$ satisfies

$$S(k) = I_k - 2\pi i A(k), \qquad A(k) \in \mathbf{B}_1, \tag{1.37}$$

$$\det S(k) = \frac{\overline{D}(k + i0)}{D(k + i0)}. \tag{1.38}$$

Remark. 1) Consider Schrödinger operators $\widetilde{H} = -\Delta + \widetilde{Q}$ on $L^2(\mathbb{R}^d), d \geqslant 1$, where $-\Delta$ is the standard Laplacian in $\mathbb{R}^d$ and $\widetilde{Q} \in C_0^\infty(\mathbb{R}^d)$ is a real potential. Define the operator $\widetilde{Y}_0(k) = |\widetilde{Q}|^{\frac12}(-\Delta - k^2)^{-1}\widetilde{Q}^{\frac12}, k \in \mathbb{C}_+$.

Firstly, if $d = 1$, then it is well known that the corresponding operator $\widetilde{Y}_0(k) \in \mathbf{B}_1$ and so the Fredholm determinant $\det(I + \widetilde{Y}_0(k))$ is well defined. Recall that the Fredholm determinant is the basic function to study trace formulas, spectral shift functions, etc. in the general framework.

Secondly, if $d \geqslant 2$, then it is well known that the corresponding operator $\widetilde{Y}_0(k)$ is not trace class and so the Fredholm determinant $\det(I+\widetilde{Y}_0(k))$ is not defined. Thus, we need an essential modification. For example, Newton [N77] defined the modified Fredholm determinant by

$$\widetilde{D}(k) = \det[(I + \widetilde{Y}_0(k)e^{-\widetilde{Y}_0(k)}], \qquad k \in \mathbb{C}_+, \quad if \quad d = 3.$$

Note that the case $d > 3$ is more complicated.

Thus, in contrast to Schrödinger operators $\widetilde{H}$ on $\mathbb{R}^d$ in the case of Schrödinger operators $H = \Delta_M + Q$ on metric graphs the operator $Y_0(k)$ is trace class and the corresponding Fredholm determinant $D(k) = \det(I + Y_0(k))$ is well define for all $k \in \mathbb{C}_+$ and for any dimension $d \geqslant 1$. Then the metric case (for any dimension $d \geqslant 1$) is very similar to the case of Schrödinger operators $\widetilde{H}$ on $\mathbb{R}^1$. This fact is very important for spectral theory of Schrödinger operators on periodic metric graphs.

2) In fact the identity (1.38) is the so-called Birman-Krein [BK62] formula for S-matrix in the case of the scattering on metric graphs.

3) Recall that if the scattering operator $S = W_+^* W_-$ is unitary on $\mathscr{H}_{ac}(H_0)$, then the operators H_0 and S commute and thus are simultaneously diagonalizable:

$$\mathscr{H}_{ac}(H_0) = P_{ac}(H_0)L^2(\Gamma) = \int_\sigma^\oplus \mathscr{H}_\lambda d\lambda, \qquad H_0 = \int_\sigma^\oplus \lambda I_\lambda d\lambda, \qquad S = \int_\sigma^\oplus S(\lambda) d\lambda; \quad (1.39)$$

here $\sigma = \sigma_{ac}(H_0)$ and I_λ is the identity in the fiber space $\mathscr{H}_\lambda$ and $S(\lambda)$ is the scattering matrix acting in $\mathscr{H}_\lambda$ for the pair H_0, H.

2. Floquet decomposition of metric Laplacians

2.1. **Definitions.** We introduce *an edge index*, which is important to study the spectrum of Schrödinger operators on periodic graphs. We fix any ν vertices of the periodic graph Γ, which are not $\mathbb{Z}^d$-equivalent to each other and denote this vertex set by V_0. We will call V_0 *a fundamental vertex set of* Γ. For any $v \in V$ the following unique representation holds true:

$$v = v_0 + [v], \qquad v_0 \in V_0, \qquad [v] \in \mathbb{Z}^d. \tag{2.1}$$

In other words, each vertex v can be obtained from a vertex $v_0 \in V_0$ by the shift by a vector $[v] \in \mathbb{Z}^d$. We will call $[v]$ *coordinates of the vertex v with respect to the fundamental vertex set V_0*. For any oriented edge $\mathbf{e} = (u, v) \in \mathcal{A}$ we define **the edge "index"** $\tau(\mathbf{e})$ as the integer vector by

$$\tau(\mathbf{e}) = [v] - [u] \in \mathbb{Z}^d, \tag{2.2}$$

where, due to (2.1), we have

$$u = u_0 + [u], \qquad v = v_0 + [v], \qquad u_0, v_0 \in V_0, \qquad [u], [v] \in \mathbb{Z}^d.$$

In general, edge indices depend on the choice of the set V_0. We note that edges connecting vertices from the fundamental vertex set V_0 have zero indices.

We define two surjections

$$\mathfrak{f}_V : V \to V_* = V/\mathbb{Z}^d, \qquad \mathfrak{f}_\mathcal{A} : \mathcal{A} \to \mathcal{A}_* = \mathcal{A}/\mathbb{Z}^d, \tag{2.3}$$

which map each element to its equivalence class.

If $\mathbf{e}$ is an oriented edge of the graph Γ, then by the definition of the fundamental graph there is an oriented edge $\mathbf{e}_* = \mathfrak{f}_{\mathcal{A}}(\mathbf{e})$ on Γ_*. For the edge $\mathbf{e}_* \in \mathcal{A}_*$ we define the edge index $\tau(\mathbf{e}_*)$ by

$$\tau(\mathbf{e}_*) = \tau(\mathbf{e}). \tag{2.4}$$

In other words, edge indices of the fundamental graph Γ_* are induced by edge indices of the periodic graph Γ. The index of a fundamental graph edge with respect to the fixed fundamental vertex set V_0 is uniquely determined by (2.4), since

$$\tau(\mathbf{e}+m) = \tau(\mathbf{e}), \qquad \forall\, (\mathbf{e}, m) \in \mathcal{A} \times \mathbb{Z}^d.$$

Edges with nonzero indices will be called **bridges**. They are important to describe the spectrum of the Laplacian. The bridges provide the connectivity of the periodic graph and the removal of all bridges disconnects the graph into infinitely many connected components. The set of all bridges of the fundamental graph Γ_* we denote by $\mathcal{B}_*$.

Example. We consider the periodic graph, shown in Fig.1. The periods of the graph a_1, a_2 are also shown in this figure. We choose and fix a fundamental vertex set $V_0 = \{v_1, v_2, v_3, v_4\}$ such that Γ_0, the subgraph of Γ induced by V_0, is connected. Here the subgraph Γ_0 of Γ induced by V_0 is a graph that consists of all vertices from V_0 and all edges of Γ that connect these vertices.



FIGURE 1. Periodic graph Γ; the fundamental vertices v_1, v_2, v_3, v_4 are black; the bridges of Γ are marked by bold.

2.2. **Floquet fiber Laplacians.** The normalized Laplacian Δ, given by (1.3), has the standard decomposition into a constant fiber direct integral (1.4), where the Floquet $\nu \times \nu$ matrix $\Delta(\vartheta) = \{\Delta_{uv}(\vartheta)\}_{u,v \in V_*}$ has the form

$$\Delta_{uv}(\vartheta) = \begin{cases} -\frac{1}{\sqrt{\varkappa_u \varkappa_v}} \sum\limits_{\mathbf{e}=(u,v)\in\mathcal{A}_*} e^{\,i\langle \tau(\mathbf{e}),\,\vartheta\rangle}, & \text{if } \ (u,v) \in \mathcal{A}_* \\ \quad 0, & \text{if } \ (u,v) \notin \mathcal{A}_* \end{cases}, \tag{2.5}$$

see Theorem 2.2 in [KS15c], where $\varkappa_v$ is the degree of the vertex v, $\langle\cdot\,,\cdot\rangle$ denotes the standard inner product in $\mathbb{R}^d$, the vector $\vartheta \in \mathbb{T}^d$ is called the quasimomentum and $\tau(\mathbf{e})$ is the edge index defined by (2.2), (2.4).

We identify an edge $\mathbf{e}_* = (u_*, v_*) \in \mathcal{E}_*$ of the fundamental graph $\Gamma_* = (V_*, \mathcal{E}_*)$ with an index $\tau(\mathbf{e}_*)$ with the edge $\mathbf{e} = (u, v + \tau(\mathbf{e}_*)) \in \mathcal{E}$ of the graph $\Gamma = (V, \mathcal{E})$, where $u, v \in V_0$ such that $u_* = \mathfrak{f}_V(u)$, $v_* = \mathfrak{f}_V(v)$. We note that $\tau(\mathbf{e}) = \tau(\mathbf{e}_*)$.

Proof of Theorem 1.1. Denote by $L^2_{com}(\Gamma)$ the set of all compactly supported functions $h \in L^2(\Gamma)$. Standard arguments (see pp. 290–291 in [RS78]) yield that $\mathscr{U}$ given by (1.11), is well defined on $L^2_{com}(\Gamma)$ and has a unique extension to a unitary operator. For $h \in L^2_{com}(\Gamma)$ the sum (1.11) is finite and, using the identity $\mathcal{E} = \big\{\mathbf{e}_* + m : (\mathbf{e}_*, m) \in \mathcal{E}_* \times \mathbb{Z}^d\big\}$, we have

$$\begin{aligned}
\|\mathscr{U}h\|^2_{\mathscr{H}} &= \int_{\mathbb{T}^d} \|(\mathscr{U}h)(\vartheta,\cdot)\|^2_{L^2(\Gamma_*)} \frac{d\vartheta}{(2\pi)^d} \\
&= \sum_{\mathbf{e}_*\in\mathcal{E}_*} \int_0^1 \int_{\mathbb{T}^d} \Big(\sum_{m\in\mathbb{Z}^d} e^{-i\langle m,\,\vartheta\rangle} h_{\mathbf{e}_*+m}(t)\Big)\Big(\sum_{m'\in\mathbb{Z}^d} e^{i\langle m',\,\vartheta\rangle} \bar{h}_{\mathbf{e}_*+m'}(t)\Big) \frac{d\vartheta}{(2\pi)^d}\, dt \\
&= \sum_{\mathbf{e}_*\in\mathcal{E}_*} \int_0^1 \sum_{m,m'\in\mathbb{Z}^d} \bigg(h_{\mathbf{e}_*+m}(t)\bar{h}_{\mathbf{e}_*+m'}(t) \int_{\mathbb{T}^d} e^{-i\langle m-m',\,\vartheta\rangle} \frac{d\vartheta}{(2\pi)^d}\bigg)\, dt \\
&= \sum_{\mathbf{e}_*\in\mathcal{E}_*} \int_0^1 \sum_{m\in\mathbb{Z}^d} \big|h_{\mathbf{e}_*+m}(t)\big|^2\, dt = \sum_{\mathbf{e}\in\mathcal{E}} \int_0^1 |h_{\mathbf{e}}(t)|^2\, dt = \|h\|^2_{L^2(\Gamma)}.
\end{aligned}$$

Thus, $\mathscr{U}$ is well defined on $L^2_{com}(\Gamma)$ and has a unique isometric extension. In order to prove that $\mathscr{U}$ is onto $\mathscr{H}$ we compute $\mathscr{U}^*$. Let $g = \big(g(\cdot,x_*)\big)_{x_*\in\Gamma_*} \in \mathscr{H}$, where $g(\cdot,x_*) : \mathbb{T}^d \to \mathbb{C}$. We define

$$(\mathscr{U}^*g)(x) = \int_{\mathbb{T}^d} e^{i\langle m,\vartheta\rangle} g(\vartheta,x_*) \frac{d\vartheta}{(2\pi)^d}\,, \qquad x = x_* + m \in \Gamma, \tag{2.6}$$

where $(x_*, m) \in \Gamma_* \times \mathbb{Z}^d$ are uniquely defined. A direct computation gives that this is indeed the formula for the adjoint of $\mathscr{U}$. Moreover, the Parseval identity for Fourier series gives

$$\|\mathscr{U}^*g\|^2_{L^2(\Gamma)} = \sum_{\mathbf{e}\in\mathcal{E}} \big\|(\mathscr{U}^*g)_{\mathbf{e}}\big\|^2_{L^2(0,1)} = \sum_{\mathbf{e}_*\in\mathcal{E}_*} \sum_{m\in\mathbb{Z}^d} \int_0^1 \big|(\mathscr{U}^*g)_{\mathbf{e}_*+m}(t)\big|^2\, dt$$

$$= \sum_{\mathbf{e}_*\in\mathcal{E}_*} \sum_{m\in\mathbb{Z}^d} \int_0^1 \bigg|\int_{\mathbb{T}^d} e^{i\langle m,\vartheta\rangle} g_{\mathbf{e}_*}(\vartheta,t) \frac{d\vartheta}{(2\pi)^d}\bigg|^2 dt$$

$$= \sum_{\mathbf{e}_*\in\mathcal{E}_*} \int_0^1 \int_{\mathbb{T}^d} \big|g_{\mathbf{e}_*}(\vartheta,t)\big|^2 \frac{d\vartheta}{(2\pi)^d}\, dt = \int_{\mathbb{T}^d} \big\|g(\vartheta,\cdot)\big\|^2_{L^2(\Gamma_*)} \frac{d\vartheta}{(2\pi)^d} = \|g\|^2_{\mathscr{H}}.$$

Further, for $h \in L^2_{com}(\Gamma)$ and $x \in \mathbf{e} \in \mathcal{E}_*$, $x \notin V_*$ we obtain

$$\begin{aligned}
(\mathscr{U}\Delta_M h)(\vartheta,x) &= \sum_{m\in\mathbb{Z}^d} e^{-i\langle m,\vartheta\rangle} (\Delta_M h)(x+m) \\
&= \sum_{m\in\mathbb{Z}^d} e^{-i\langle m,\vartheta\rangle} h''_{\mathbf{e}+m}(x+m) = \Delta_M(\vartheta)(\mathscr{U}h)(\vartheta,x).
\end{aligned} \tag{2.7}$$

Let $v\in V_*$ and $\mathbf{e}_1,\mathbf{e}_2,\mathbf{e}\in I_*(v)$. Denote by $\delta_1=\delta(\mathbf{e}_1,v)$, $\delta_2=\delta(\mathbf{e}_2,v)$ and $\delta=\delta(\mathbf{e},v)$, where δ is defined by (1.2). Then, using (1.1), we have

$$\begin{aligned}
e^{-i\delta_1\langle\tau(\mathbf{e}_1),\vartheta\rangle}(\mathscr{U}h)_{\mathbf{e}_1}(\vartheta,\delta_1)&=e^{-i\delta_1\langle\tau(\mathbf{e}_1),\vartheta\rangle}\sum_{m\in\mathbb{Z}^d}e^{-i\langle m,\vartheta\rangle}h_{\mathbf{e}_1+m}(\delta_1)\\
&=e^{-i\delta_2\langle\tau(\mathbf{e}_2),\vartheta\rangle}\sum_{m\in\mathbb{Z}^d}e^{-i\langle m+\delta_1\tau(\mathbf{e}_1)-\delta_2\tau(\mathbf{e}_2),\vartheta\rangle}h_{\mathbf{e}_1+m}(\delta_1)\\
&=e^{-i\delta_2\langle\tau(\mathbf{e}_2),\vartheta\rangle}\sum_{m\in\mathbb{Z}^d}e^{-i\langle m+\delta_1\tau(\mathbf{e}_1)-\delta_2\tau(\mathbf{e}_2),\vartheta\rangle}h_{\mathbf{e}_2+m+\delta_1\tau(\mathbf{e}_1)-\delta_2\tau(\mathbf{e}_2)}(\delta_2)\\
&=e^{-i\delta_2\langle\tau(\mathbf{e}_2),\vartheta\rangle}\sum_{m\in\mathbb{Z}^d}e^{-i\langle m,\vartheta\rangle}h_{\mathbf{e}_2+m}(\delta_2)=e^{-i\delta_2\langle\tau(\mathbf{e}_2),\vartheta\rangle}(\mathscr{U}h)_{\mathbf{e}_2}(\vartheta,\delta_2),
\end{aligned}\tag{2.8}$$

and

$$\begin{aligned}
\sum_{\mathbf{e}\in I_*(v)}(-1)^{\delta}e^{-i\delta\langle\tau(\mathbf{e}),\,\vartheta\rangle}\,(\mathscr{U}h)'_{\mathbf{e}}(\vartheta,\delta)&=\sum_{\mathbf{e}\in I_*(v)}(-1)^{\delta}e^{-i\delta\langle\tau(\mathbf{e}),\,\vartheta\rangle}\sum_{m\in\mathbb{Z}^d}e^{-i\langle m,\vartheta\rangle}h'_{\mathbf{e}+m}(\delta)\\
=\sum_{\mathbf{e}\in I_*(v)}(-1)^{\delta}\sum_{m\in\mathbb{Z}^d}e^{-i\langle m+\delta\tau(\mathbf{e}),\vartheta\rangle}h'_{\mathbf{e}+m}(\delta)&=\sum_{\mathbf{e}\in I_*(v)}(-1)^{\delta}\sum_{m\in\mathbb{Z}^d}e^{-i\langle m,\vartheta\rangle}h'_{\mathbf{e}+m-\delta\tau(\mathbf{e})}(\delta)\\
=\sum_{m\in\mathbb{Z}^d}e^{-i\langle m,\vartheta\rangle}\sum_{\mathbf{e}\in I_*(v)}(-1)^{\delta}h'_{\mathbf{e}+m-\delta\tau(\mathbf{e})}(\delta)&=\sum_{m\in\mathbb{Z}^d}e^{-i\langle m,\vartheta\rangle}\sum_{\mathbf{e}\in I(v+m)}(-1)^{\delta}h'_{\mathbf{e}+m}(\delta)=0.
\end{aligned}\tag{2.9}$$

The identities (2.7) – (2.9) yield (1.10). ■

Corollary 2.1. *The Laplacian Δ_M on $L^2(\Gamma)$ has a decomposition into a constant fiber direct integral of the form (1.10), where the fiber operator $\widetilde{\Delta}_M(\vartheta)$ on $L^2(\Gamma_*)$ is defined by*

$$(\widetilde{\Delta}_M(\vartheta)y\,)_{\mathbf{e}}=-\big(\tfrac{d}{dt}+i\,\langle\tau(\mathbf{e}),\,\vartheta\rangle\big)^2y_{\mathbf{e}},\qquad y=(y_{\mathbf{e}})_{\mathbf{e}\in\mathcal{E}_*},\tag{2.10}$$

where $(y''_{\mathbf{e}})_{\mathbf{e}\in\mathcal{E}_*}\in L^2(\Gamma_*)$, y is continuous on Γ_* and satisfies the boundary conditions:

$$\sum_{\mathbf{e}\in I_*(v)}(-1)^{\delta(\mathbf{e},v)}\big(\tfrac{d}{dt}+i\,\langle\tau(\mathbf{e}),\vartheta\rangle\big)\,y_{\mathbf{e}}\big(\delta(\mathbf{e},v)\big)=0\qquad\forall v\in V_*,\tag{2.11}$$

or, that is the same,

$$\begin{aligned}
\sum_{\mathbf{e}\in I_*(v)}(-1)^{\delta(\mathbf{e},v)}y'_{\mathbf{e}}\big(\delta(\mathbf{e},v)\big)&=y(v)\,i\sum_{\mathbf{e}\in I_*(v)}(-1)^{\delta(\mathbf{e},v)}\langle\tau(\mathbf{e}),\vartheta\rangle,\quad\forall v\in V_*,\\
y(v)&=y_{\mathbf{e}}\big(\delta(\mathbf{e},v)\big),\quad\forall\,\mathbf{e}\in I_*(v).
\end{aligned}\tag{2.12}$$

Proof. After the gauge transformation $\widetilde{y}_{\mathbf{e}}(t)=e^{-it\langle\tau(\mathbf{e}),\,\vartheta\rangle}y_{\mathbf{e}}(t)$, for all $\mathbf{e}\in\mathcal{E}_*$, identities (1.12) – (1.14) take the form (2.10) – (2.11). ■

3. Eigenfunctions of metric Laplacians

In order to prove Theorem 1.2 we need to choose the fundamental vertex set V_0 by a specific way. Let $T=(V_T,\mathcal{E}_T)$ be a subgraph of the periodic graph Γ satisfying the following conditions:

1) *T is a tree, i.e., a connected graph without cycles;*

2) *V_T is a fundamental vertex set, i.e., V_T consists of ν vertices of Γ, which are not $\mathbb{Z}^d$-equivalent to each other. Recall that ν is the number of vertices of Γ_*.*

Remark. 1) We need to note that such a graph T always exists, since the periodic graph is connected, and T is not unique.

2) The graph $T_*=T/\mathbb{Z}^d$ is a spanning tree of the fundamental graph Γ_*, i.e., $T_*=(V_*,\mathcal{E}_{T_*})$ is a subgraph of Γ_*, which has $\nu-1$ edges and contains no cycles.

From now on we assume that the fundamental vertex set V_0 coincides with the vertex set V_T. Then, by the definition of the edge index, **all edges of the spanning tree T_* have zero indices**.

The following lemma holds true (see Lemma 2.3 in [KS15c]).

Lemma 3.1. *Let $\mathbf{e}\in\mathcal{B}_*$ be a bridge of the fundamental graph Γ_* with the index $\tau(\mathbf{e})$. Then*

i) There exists a unique cycle in Γ_*, containing only $\mathbf{e}$ and edges of T_*.

ii) The length of this cycle, i.e., the number of its edges, is not more than ν and the sum of all indices of the cycle edges is $\tau(\mathbf{e})$.

Proof of Theorem 1.2. i)-iii) We construct eigenfunctions $\Psi_j^0=\big(\Psi_{j,\mathbf{e}}^0\big)_{\mathbf{e}\in\mathcal{E}_*}$ of the operator $\Delta_M(\vartheta)$ corresponding to the eigenvalue $(\pi j)^2$. These eigenfunctions have the form

$$\Psi_{j,\mathbf{e}}^0(\vartheta,t)=\Psi_{j,\mathbf{e}}^0(\vartheta,0)\,\cos(\pi jt)+{\Psi_{j,\mathbf{e}}^0}'(\vartheta,0)\,\tfrac{\sin(\pi jt)}{\pi j}\,,\qquad \forall\, t\in[0,1],\tag{3.1}$$

which yields

$$\Psi_{j,\mathbf{e}}^0(\vartheta,1)=(-1)^j\,\Psi_{j,\mathbf{e}}^0(\vartheta,0),\qquad {\Psi_{j,\mathbf{e}}^0}'(\vartheta,1)=(-1)^j\,{\Psi_{j,\mathbf{e}}^0}'(\vartheta,0).\tag{3.2}$$

Since Γ is connected, on Γ, and consequently, on the fundamental graph Γ_*, there exist d bridges $\widetilde{\mathbf{e}}_1,\ldots,\widetilde{\mathbf{e}}_d\in\mathcal{B}_*$ with linearly independent indices $\tau(\widetilde{\mathbf{e}}_1),\ldots,\tau(\widetilde{\mathbf{e}}_d)\in\mathbb{Z}^d$. Lemma 3.1 gives that on Γ_* there exist cycles W_s of length l_s, $s\in\mathbb{N}_d$, each of which contains only $\widetilde{\mathbf{e}}_s$ and edges of T_*. Recall that all edges of T_* have zero indices. Then the quasi-periodic condition (1.13) and the first identity in (3.2) yield

$$\Psi_{j,\widetilde{\mathbf{e}}_s}^0(\vartheta,0)\big((-1)^{jl_s}\,e^{i\langle\tau(\widetilde{\mathbf{e}}_s),\,\vartheta\rangle}-1\big)=0,\qquad \forall\,(\vartheta,s)\in\mathbb{T}^d\times\mathbb{N}_d.\tag{3.3}$$

Since the indices $\tau(\widetilde{\mathbf{e}}_1),\ldots,\tau(\widetilde{\mathbf{e}}_d)$ are linearly independent, from (3.3) it follows that $\Psi_{j,\widetilde{\mathbf{e}}_s}^0(\vartheta,0)=0$, $s\in\mathbb{N}_d$, for all $\vartheta\in\mathbb{T}^d\setminus\{0\}$. Due to the connectivity of the fundamental graph, the quasi-periodic condition (1.13) and the first identity in (3.2) give $\Psi_{j,\mathbf{e}}^0(\vartheta,0)=0$ for all $\mathbf{e}\in\mathcal{E}_*$. Thus, the eigenfunctions (3.1) take the form

$$\Psi_{j,\mathbf{e}}^0(\vartheta,t)={\Psi_{j,\mathbf{e}}^0}'(\vartheta,0)\,\tfrac{\sin(\pi jt)}{\pi j}\,,\qquad \forall\,\mathbf{e}\in\mathcal{E}_*.\tag{3.4}$$

These functions must satisfy the quasi-periodic condition (1.14), which, using the second identity in (3.2), can be rewritten in the form

$$\sum_{\mathbf{e}=(v,u)\in\mathcal{E}_*}{\Psi_{j,\mathbf{e}}^0}'(\vartheta,0)-(-1)^j\sum_{\mathbf{e}=(u,v)\in\mathcal{E}_*}e^{-i\langle\tau(\mathbf{e}),\,\vartheta\rangle}\,{\Psi_{j,\mathbf{e}}^0}'(\vartheta,0)=0,\qquad \forall v\in V_*.\tag{3.5}$$

This is a homogeneous system of $\nu=\#V_*$ linear equations with $\nu_*=\#\mathcal{E}_*$ variables $\big({\Psi_{j,\mathbf{e}}^0}'(\vartheta,0)\big)_{\mathbf{e}\in\mathcal{E}_*}$. The $\nu\times\nu_*$ coefficient matrix $D_j(\vartheta)=\{D_{v,\mathbf{e}}^{(j)}(\vartheta)\}_{\substack{v\in V_*\\ \mathbf{e}\in\mathcal{E}_*}}$ of the system is given by

$$D_{v,\mathbf{e}}^{(j)}(\vartheta)=\begin{cases}(-1)^{j+1}e^{-i\langle\tau(\mathbf{e}),\,\vartheta\rangle}, & \text{if } v \text{ is the terminal vertex of } \mathbf{e}\\ 1, & \text{if } v \text{ is the initial vertex of } \mathbf{e}\\ 1-(-1)^je^{-i\langle\tau(\mathbf{e}),\,\vartheta\rangle} & \text{if } \mathbf{e} \text{ is a loop in the vertex } v\\ 0, & \text{otherwise}\end{cases}\ .\tag{3.6}$$

The number of linearly independent solutions of the system (3.5) is $\big(\nu_* - \operatorname{rank} D_j(\vartheta)\big)$. We will show that $\operatorname{rank} D_j(\vartheta) = \nu$ for all $\vartheta \in \mathbb{T}^d \setminus \{0\}$.

Let $D^{(j)}_{v\boldsymbol{\cdot}}(\vartheta)$, $v \in V_*$, denote the row of the matrix $D_j(\vartheta)$, corresponding to the vertex v. In order to show that the rows of this matrix are linearly independent, we consider their linear combination with coefficients α_v, $v \in V_*$:

$$\sum_{v \in V_*} \alpha_v \, D^{(j)}_{v\boldsymbol{\cdot}}(\vartheta) = 0. \tag{3.7}$$

From this, using the form (3.6) of the matrix $D_j(\vartheta)$, we obtain

$$\alpha_u = \alpha_v (-1)^j e^{-i\langle \tau(\mathbf{e}),\, \vartheta\rangle}, \qquad \forall\, \mathbf{e} = (u,v) \in \mathcal{E}_*. \tag{3.8}$$

For each vertex v of the cycle W_s the identity (3.8) gives

$$\alpha_v \big((-1)^{jl_s} e^{-i\langle \tau(\widetilde{\mathbf{e}}_s),\, \vartheta\rangle} - 1 \big) = 0, \qquad \forall\, (\vartheta, s) \in \mathbb{T}^d \times \mathbb{N}_d. \tag{3.9}$$

Since $\vartheta \neq 0$ and $\tau(\widetilde{\mathbf{e}}_1), \ldots, \tau(\widetilde{\mathbf{e}}_d)$ are linearly independent, we conclude that $\alpha_v = 0$. Due to the connectivity of Γ_* and the identity (3.8), all coefficients $\alpha_v = 0$ in (3.7). Thus, for all $\vartheta \in \mathbb{T}^d \setminus \{0\}$ the rows of the matrix $D_j(\vartheta)$ are linearly independent and $\operatorname{rank} D_j(\vartheta) = \nu$. The number $\nu_* - \nu$ of the linearly independent solutions of the system (3.5) is equal to the multiplicity of each eigenvalue $(\pi j)^2$.

From (3.4) it follows that

$$\|\Psi^0_j(\vartheta, \cdot)\|^2_{L^2(\Gamma_*)} = \frac{1}{2(\pi j)^2} \sum_{\mathbf{e} \in \mathcal{E}_*} \big| {\Psi^0_{j,\mathbf{e}}}'(\vartheta, 0) \big|^2. \tag{3.10}$$

Choosing the normalized solutions $\big(X_{s,j,\mathbf{e}}(\vartheta)\big)_{\mathbf{e} \in \mathcal{E}_*}$, $s \in \mathbb{N}_{\nu_* - \nu}$, of the system (3.5) or, equivalently, of the system (1.18), we obtain that the normalized eigenfunctions $\Psi^0_{s,j} = \big(\Psi^0_{s,j,\mathbf{e}}\big)_{\mathbf{e} \in \mathcal{E}_*}$, $s \in \mathbb{N}_{\nu_* - \nu}$, have the form (1.17).

The identity $X_{s,j,\mathbf{e}}(\vartheta) = X_{s,j+2,\mathbf{e}}(\vartheta)$, $(s, \mathbf{e}) \in \mathbb{N}_{\nu_* - \nu} \times \mathcal{E}_*$, follows from the form of the system (1.18). The inequality (1.19) is a direct consequence of (1.17). ■

Remark. The Floquet operator $\Delta_M(0)$ is the metric Laplacian on the fundamental graph Γ_*. It is known (see [B85], [C97], [LP08]), that

i) the normalized eigenfunctions $\Psi^0_{s,j} = \big(\Psi^0_{s,j,\mathbf{e}}\big)_{\mathbf{e} \in \mathcal{E}_*}$, $s \in \mathbb{N}_{\nu_* - \nu + 1}$ of the operator $\Delta_M(0)$ corresponding to the eigenvalue $(\pi j)^2$, j is even, have the form (1.17) as $\vartheta = 0$, where $\big(X_{s,j,\mathbf{e}}(0)\big)_{\mathbf{e} \in \mathcal{E}_*}$ is a normalized solution of the system of ν equations

$$\sum_{\substack{\mathbf{e} \in \mathcal{E}_*, \\ \text{starting at } v}} x_{\mathbf{e}} - \sum_{\substack{\mathbf{e} \in \mathcal{E}_*, \\ \text{ending at } v}} x_{\mathbf{e}} = 0, \qquad \forall v \in V_*, \tag{3.11}$$

one equation for each $v \in V_*$. This system has the rank $\nu - 1$ and thus there exist $\nu_* - \nu + 1$ linearly independent normalized solutions $\big(X_{s,j,\mathbf{e}}(0)\big)_{\mathbf{e} \in \mathcal{E}_*}$, $s \in \mathbb{N}_{\nu_* - \nu + 1}$. In this case there exists an additional normalized eigenfunction $\Psi^0_j = \big(\Psi^0_{j,\mathbf{e}}\big)_{\mathbf{e} \in \mathcal{E}_*}$, having the form

$$\Psi^0_{j,\mathbf{e}}(0, t) = \sqrt{\frac{2}{\nu_*}} \cos(\pi j t). \tag{3.12}$$

This additional eigenfunction is related to the eigenfunction of the discrete Laplacian $\Delta(0)$ with the eigenvalue -1.

ii) the normalized eigenfunctions $\Psi^0_{s,j}=\big(\Psi^0_{s,j,\mathbf{e}}\big)_{\mathbf{e}\in\mathcal{E}_*}$ corresponding to the eigenvalue $(\pi j)^2$, j is odd, also have the form (1.17) as $\vartheta=0$, where $\big(X_{s,j,\mathbf{e}}(0)\big)_{\mathbf{e}\in\mathcal{E}_*}$ is a normalized solution of the system of ν equations

$$\sum_{\substack{\mathbf{e}\in\mathcal{E}_*,\\ \text{starting at } v}} x_{\mathbf{e}}+\sum_{\substack{\mathbf{e}\in\mathcal{E}_*,\\ \text{ending at } v}} x_{\mathbf{e}}=0,\qquad \forall v\in V_*. \tag{3.13}$$

If Γ_* is bipartite, this system has the rank $\nu-1$. If Γ_* is not bipartite, this system has the rank ν. For a bipartite graph there exists an additional normalized eigenfunction $\Psi^0_j=\big(\Psi^0_{j,\mathbf{e}}\big)_{\mathbf{e}\in\mathcal{E}_*}$, having the form

$$\Psi^0_{j,\mathbf{e}}(0,t)=\pm\sqrt{\frac{2}{\nu_*}}\cos(\pi jt), \tag{3.14}$$

where the sign is chosen such that the eigenfunction has alternative sign in adjacent vertices. This additional eigenfunction is related to the eigenfunction of the discrete Laplacian $\Delta(0)$ with the eigenvalue 1.

In the following proposition we describe the connection between eigenfunctions and corresponding eigenvalues (except for the Dirichlet eigenvalues σ_D) of the Floquet operators for metric and discrete Laplacians.

Proposition 3.2. *Let $\vartheta\in\mathbb{T}^d$. Then the following statements hold true.*

i) If $\Psi(\vartheta)=\big(\Psi_{\mathbf{e}}(\vartheta,t)\big)_{\mathbf{e}\in\mathcal{E}_*}\in L^2(\Gamma_*)$ is an eigenfunction of the operator $\Delta_M(\vartheta)$ with an eigenvalue $E(\vartheta)\notin\sigma_D$, then the function $\psi(\vartheta,\cdot)\in\ell^2(V_*)$, defined by

$$\psi(\vartheta,v)=\sqrt{\varkappa_v}\,e^{-i\delta(\mathbf{e},v)\langle\tau(\mathbf{e}),\,\vartheta\rangle}\Psi_{\mathbf{e}}(\vartheta,\delta(\mathbf{e},v)), \tag{3.15}$$

is an eigenfunction of the operator $\Delta(\vartheta)$ with the eigenvalue $\lambda(\vartheta)=-\cos\sqrt{E(\vartheta)}$.

ii) Conversely, if $\psi(\vartheta,\cdot)\in\ell^2(V_*)$ is an eigenfunction of $\Delta(\vartheta)$ with an eigenvalue $\lambda(\vartheta)\in(-1,1)$, then for each $j\in\mathbb{N}$ the function $\Psi_j(\vartheta)=\big(\Psi_{\mathbf{e},j}(\vartheta,t)\big)_{\mathbf{e}\in\mathcal{E}_*}\in L^2(\Gamma_*)$, defined by

$$\begin{aligned}&\Psi_{\mathbf{e},j}(\vartheta,t)=\tfrac{1}{\sin z_j(\vartheta)}\Big(\tfrac{\psi(\vartheta,u)}{\sqrt{\varkappa_u}}\sin(z_j(\vartheta)(1-t))+\tfrac{\psi(\vartheta,v)}{\sqrt{\varkappa_v}}e^{i\langle\tau(\mathbf{e}),\,\vartheta\rangle}\sin(z_j(\vartheta)\,t)\Big),\qquad \mathbf{e}=(u,v),\\ &z_j(\vartheta)=\begin{cases}z(\vartheta)+\pi j, & j\ \text{is even}\\ (\pi-z(\vartheta))+\pi j, & j\ \text{is odd}\end{cases},\qquad z(\vartheta)=\arccos(-\lambda(\vartheta)),\end{aligned} \tag{3.16}$$

is an eigenfunction of $\Delta_M(\vartheta)$ with the eigenvalue $E_j(\vartheta)=z_j^2(\vartheta)$.

Proof. i) Let $\Psi(\vartheta)$ be an eigenfunction of the operator $\Delta_M(\vartheta)$ with an eigenvalue $E\notin\sigma_D$, $E\neq0$, $E\equiv E(\vartheta)$. The proof for the case $E=0$ is similar. Then

$$\Psi_{\mathbf{e}}(t)=\Psi_{\mathbf{e}}(0)\cos(zt)+\Psi'_{\mathbf{e}}(0)\,\tfrac{\sin(zt)}{z},\qquad z=\sqrt{E},\qquad \Psi_{\mathbf{e}}(t)\equiv\Psi_{\mathbf{e}}(\vartheta,t), \tag{3.17}$$

which yields

$$\Psi_{\mathbf{e}}(1)=\Psi_{\mathbf{e}}(0)\cos z+\Psi'_{\mathbf{e}}(0)\,\tfrac{\sin z}{z},\qquad \Psi'_{\mathbf{e}}(1)=-\Psi_{\mathbf{e}}(0)z\sin z+\Psi'_{\mathbf{e}}(0)\cos z. \tag{3.18}$$

From (3.18) we obtain

$$\Psi'_{\mathbf{e}}(0)=\tfrac{z}{\sin z}\big(\Psi_{\mathbf{e}}(1)-\Psi_{\mathbf{e}}(0)\cos z\big),\qquad \Psi'_{\mathbf{e}}(1)=\tfrac{z}{\sin z}\big(\Psi_{\mathbf{e}}(1)\cos z-\Psi_{\mathbf{e}}(0)\big). \tag{3.19}$$

Using the formulas (3.19), we rewrite the left-hand side of the condition (1.14) in the form:

$$\begin{aligned}\sum_{\mathbf{e}\in I_*(v)}(-1)^{\delta(\mathbf{e},v)}e^{-i\delta(\mathbf{e},v)\langle\tau(\mathbf{e}),\,\vartheta\rangle}\,\Psi_{\mathbf{e}}'\big(\delta(\mathbf{e},v)\big)&=\sum_{\mathbf{e}=(v,u)\in\mathcal{E}_*}\Psi_{\mathbf{e}}'(0)-\sum_{\mathbf{e}=(u,v)\in\mathcal{E}_*}e^{-i\langle\tau(\mathbf{e}),\,\vartheta\rangle}\,\Psi_{\mathbf{e}}'(1)\\=\sum_{\mathbf{e}=(v,u)\in\mathcal{E}_*}\tfrac{z}{\sin z}\big(\Psi_{\mathbf{e}}(1)-\Psi_{\mathbf{e}}(0)\cos z\big)&-\sum_{\mathbf{e}=(u,v)\in\mathcal{E}_*}e^{-i\langle\tau(\mathbf{e}),\,\vartheta\rangle}\,\tfrac{z}{\sin z}\big(\Psi_{\mathbf{e}}(1)\cos z-\Psi_{\mathbf{e}}(0)\big),\end{aligned}\tag{3.20}$$

for all $v\in V_*$. Thus, after dividing by $\frac{z}{\sin z}$, the condition (1.14) takes the form

$$\sum_{\mathbf{e}=(v,u)\in\mathcal{E}_*}\big(\Psi_{\mathbf{e}}(1)-\Psi_{\mathbf{e}}(0)\,\cos z\big)-\sum_{\mathbf{e}=(u,v)\in\mathcal{E}_*}e^{-i\langle\tau(\mathbf{e}),\,\vartheta\rangle}\,\big(\Psi_{\mathbf{e}}(1)\cos z-\Psi_{\mathbf{e}}(0)\big)=0,\tag{3.21}$$

for all $v\in V_*$. Using the formula (1.13) and introducing the notation

$$\widetilde{\Psi}(v)=e^{-i\delta(\mathbf{e},v)\langle\tau(\mathbf{e}),\,\vartheta\rangle}\Psi_{\mathbf{e}}(\delta(\mathbf{e},v)),\qquad\forall\,(v,\mathbf{e})\in V_*\times I_*(v),\tag{3.22}$$

we write the left-hand side of the identity (3.21) in the form

$$\begin{aligned}-\widetilde{\Psi}(v)\varkappa_v\cos z&+\sum_{\mathbf{e}=(v,u)\in\mathcal{E}_*}\Psi_{\mathbf{e}}(1)+\sum_{\mathbf{e}=(u,v)\in\mathcal{E}_*}e^{-i\langle\tau(\mathbf{e}),\,\vartheta\rangle}\,\Psi_{\mathbf{e}}(0)\\=-\widetilde{\Psi}(v)\varkappa_v\cos z&+\sum_{\mathbf{e}=(v,u)\in\mathcal{E}_*}e^{i\langle\tau(\mathbf{e}),\,\vartheta\rangle}\,\widetilde{\Psi}(u)+\sum_{\mathbf{e}=(u,v)\in\mathcal{E}_*}e^{-i\langle\tau(\mathbf{e}),\,\vartheta\rangle}\,\widetilde{\Psi}(u)\\&=-\widetilde{\Psi}(v)\varkappa_v\cos z+\sum_{\mathbf{e}=(v,u)\in\mathcal{A}_*}e^{i\langle\tau(\mathbf{e}),\,\vartheta\rangle}\,\widetilde{\Psi}(u).\end{aligned}\tag{3.23}$$

Thus, the condition (3.21) takes the form

$$\widetilde{\Psi}(v)\cos z=\frac{1}{\varkappa_v}\sum_{\mathbf{e}=(v,u)\in\mathcal{A}_*}e^{i\langle\tau(\mathbf{e}),\,\vartheta\rangle}\,\widetilde{\Psi}(u),\tag{3.24}$$

or, using the formulas (3.15), (3.22) and (2.5),

$$-\,\psi(\vartheta,v)\cos z=-\tfrac{1}{\sqrt{\varkappa_v}}\sum_{\mathbf{e}=(v,u)\in\mathcal{A}_*}\tfrac{1}{\sqrt{\varkappa_u}}\,e^{i\langle\tau(\mathbf{e}),\,\vartheta\rangle}\,\psi(\vartheta,u)\equiv\big(\Delta(\vartheta)\psi(\vartheta,\cdot)\big)(v).\tag{3.25}$$

Thus, $\psi(\vartheta,\cdot)$ is an eigenfunction of the operator $\Delta(\vartheta)$ with the eigenvalue $(-\cos z)$.

ii) Conversely, let $\psi(\vartheta,\cdot)$ be an eigenfunction of the operator $\Delta(\vartheta)$ with the eigenvalue $\lambda(\vartheta)\in(-1,1)$. For the function $\Psi_j(\vartheta)=\big(\Psi_{\mathbf{e},j}(\vartheta,t)\big)_{\mathbf{e}\in\mathcal{E}_*}$ defined by (3.16) we have

$$\Psi_{\mathbf{e},j}'(t)=\tfrac{z_j}{\sin z_j}\Big(-f_u\cos(z_j(1-t))+f_ve^{i\langle\tau(\mathbf{e}),\,\vartheta\rangle}\,\cos(z_jt)\Big),\tag{3.26}$$

$$\Psi_{\mathbf{e},j}''(t)=-\tfrac{z_j^2}{\sin z_j}\Big(f_u\sin(z_j(1-t))+f_ve^{i\langle\tau(\mathbf{e}),\,\vartheta\rangle}\,\sin(z_jt)\Big),\tag{3.27}$$

where for the shortness

$$\Psi_{\mathbf{e},j}(t)=\Psi_{\mathbf{e},j}(\vartheta,t),\qquad z_j=z_j(\vartheta),\qquad f_u=\frac{\psi(\vartheta,u)}{\sqrt{\varkappa_u}}\,.$$

Thus, on each edge $\mathbf{e}\in\mathcal{E}_*$ the function $\Psi_j(\vartheta)$ satisfies the equation $-\Psi_{\mathbf{e},j}''(t)=z_j^2\Psi_{\mathbf{e},j}(t)$. From (3.16) we obtain that

$$\Psi_{\mathbf{e},j}(0)=f_u,\qquad\Psi_{\mathbf{e},j}(1)=f_ve^{i\langle\tau(\mathbf{e}),\,\vartheta\rangle}\,,\qquad\mathbf{e}=(u,v),\tag{3.28}$$

which yields

$$e^{-i\delta(\mathbf{e},v)\langle\tau(\mathbf{e}),\,\vartheta\rangle}\Psi_{\mathbf{e},j}(\delta(\mathbf{e},v))=f_v,\qquad\forall\,v\in V_*,\qquad\forall\,\mathbf{e}\in I_*(v),\tag{3.29}$$

i.e., the condition (1.13) holds true. Similarly, from (3.26) we have

$$\begin{aligned} \Psi'_{\mathbf{e},j}(0) &= \tfrac{z_j}{\sin z_j}\Big(-f_u\cos z_j + f_v e^{i\langle\tau(\mathbf{e}),\,\vartheta\rangle}\Big),\\ \Psi'_{\mathbf{e},j}(1) &= \tfrac{z_j}{\sin z_j}\Big(-f_u + f_v\, e^{i\langle\tau(\mathbf{e}),\,\vartheta\rangle}\cos z_j\Big), \qquad \mathbf{e}=(u,v). \end{aligned} \tag{3.30}$$

Using (3.30), the left-hand side of the condition (1.14) takes the form

$$\begin{aligned} &\sum_{\mathbf{e}\in I_*(v)}(-1)^{\delta(\mathbf{e},v)}e^{-i\delta(\mathbf{e},v)\langle\tau(\mathbf{e}),\,\vartheta\rangle}\,\Psi'_{\mathbf{e}}\big(\delta(\mathbf{e},v)\big) = \sum_{\mathbf{e}=(v,u)\in\mathcal{E}_*}\Psi'_{\mathbf{e}}(0) - \sum_{\mathbf{e}=(u,v)\in\mathcal{E}_*}e^{-i\langle\tau(\mathbf{e}),\,\vartheta\rangle}\,\Psi'_{\mathbf{e}}(1)\\ &= \sum_{\mathbf{e}=(v,u)\in\mathcal{E}_*}\tfrac{z_j}{\sin z_j}\Big(-f_v\cos z_j + f_u e^{i\langle\tau(\mathbf{e}),\,\vartheta\rangle}\Big) + \sum_{\mathbf{e}=(u,v)\in\mathcal{E}_*}\tfrac{z_j}{\sin z_j}\Big(f_u e^{-i\langle\tau(\mathbf{e}),\,\vartheta\rangle} - f_v\cos z_j\Big)\\ &= -\tfrac{z_j\cos z_j}{\sin z_j}\varkappa_v f_v + \tfrac{z_j}{\sin z_j}\sum_{\mathbf{e}=(v,u)\in\mathcal{A}_*} f_u\, e^{i\langle\tau(\mathbf{e}),\,\vartheta\rangle}, \qquad \forall\, v\in V_*. \end{aligned}$$

This, the definition (2.5) of the operator $\Delta(\vartheta)$ and the fact that $\psi(\vartheta,\cdot)$ is an eigenfunction of $\Delta(\vartheta)$ with the eigenvalue $\lambda(\vartheta)=-\cos z_j$ yield

$$\sum_{\mathbf{e}\in I_*(v)}(-1)^{\delta(\mathbf{e},v)}e^{-i\delta(\mathbf{e},v)\langle\tau(\mathbf{e}),\,\vartheta\rangle}\,\Psi'_{\mathbf{e}}\big(\delta(\mathbf{e},v)\big) = -\tfrac{z_j\sqrt{\varkappa_v}}{\sin z_j}\Big(\psi(\vartheta,v)\cos z_j + \big(\Delta(\vartheta)\psi(\vartheta,\cdot)\big)(v)\Big) = 0$$

for all $v\in V_*$, i.e., the condition (1.14) also holds true. Thus, $\Psi_j(\vartheta)$ is an eigenfunction of the operator $\Delta_M(\vartheta)$ with an eigenvalue z_j^2. ■

Proof of Theorem 1.3. For all $\vartheta\in\mathbb{T}^d\setminus\{0\}$ the Floquet matrix $\Delta(\vartheta)$ has ν real eigenvalues $-1<\lambda_1(\vartheta)\leqslant\ldots\leqslant\lambda_\nu(\vartheta)<1$. Due to Proposition 3.2 all eigenvalues $E_{n,j}(\vartheta)=z_{n,j}^2(\vartheta)$ of the operator $\Delta_{MV}(\vartheta)$ are given by (1.22) and the eigenfunction $\Psi_{n,j}(\vartheta)$ corresponding to the eigenvalue $E_{n,j}(\vartheta)$ (defined up to a constant factor) has the form (1.23).

Now we will show (1.24). Let for the shortness

$$c_{n,j}=\cos z_{n,j}(\vartheta),\qquad s_{n,j}=\sin z_{n,j}(\vartheta),\qquad z_{n,j}=z_{n,j}(\vartheta),\quad f_n(u)=\frac{\psi_n(\vartheta,u)}{\sqrt{\varkappa_u}}\,.$$

Direct integration yields

$$\begin{aligned} \|\Psi_{n,j}(\vartheta)\|^2_{L^2(\Gamma_*)} &= \sum_{\mathbf{e}\in\mathcal{E}_*}\int_0^1\big|\Psi_{n,j,\mathbf{e}}(\vartheta,t)\big|^2dt\\ = \frac{1}{z_{n,j}s_{n,j}^2}&\sum_{\mathbf{e}=(u,v)\in\mathcal{E}_*}\bigg[\big(z_{n,j}-c_{n,j}s_{n,j}\big)\Big(|f_n(u)|^2+|f_n(v)|^2\Big)\\ +\big(s_{n,j}-z_{n,j}c_{n,j}\big)&\big(e^{-i\langle\tau(\mathbf{e}),\vartheta\rangle}f_n(u)\overline{f}_n(v)+e^{i\langle\tau(\mathbf{e}),\vartheta\rangle}\overline{f}_n(u)f_n(v)\big)\bigg]. \end{aligned} \tag{3.31}$$

The definition (2.5) of the operator $\Delta(\vartheta)$ and the fact that $\psi_n(\vartheta,\cdot)$ is an eigenfunction of $\Delta(\vartheta)$ corresponding to $\lambda_n(\vartheta)$ yield

$$
\begin{aligned}
&\sum_{\mathbf{e}=(u,v)\in\mathcal{E}_*}\big(e^{-i\langle\tau(\mathbf{e}),\vartheta\rangle}f_n(u)\overline{f}_n(v)+e^{i\langle\tau(\mathbf{e}),\vartheta\rangle}\overline{f}_n(u)f_n(v)\big)\\
&=\sum_{\mathbf{e}=(u,v)\in\mathcal{A}_*}e^{-i\langle\tau(\mathbf{e}),\vartheta\rangle}f_n(u)\overline{f}_n(v)=\sum_{u\in V_*}f_n(u)\sum_{\mathbf{e}=(u,v)\in\mathcal{A}_*}e^{-i\langle\tau(\mathbf{e}),\vartheta\rangle}\overline{f}_n(v)\\
&=-\lambda_n(\vartheta)\sum_{u\in V_*}\psi_n(\vartheta,u)\bar{\psi}_n(\vartheta,u)=c_{n,j}\|\psi_n(\vartheta,\cdot)\|^2_{\ell^2(V_*)}.
\end{aligned}
\tag{3.32}
$$

Substituting the identities

$$
\sum_{\mathbf{e}=(u,v)\in\mathcal{E}_*}\Big(|f_n(u)|^2+|f_n(v)|^2\Big)=\sum_{v\in V_*}|\psi_n(\vartheta,v)|^2=\|\psi_n(\vartheta,\cdot)\|^2_{\ell^2(V_*)}
\tag{3.33}
$$

and (3.32) into (3.31), we obtain

$$
\|\Psi_{n,j}(\vartheta)\|^2_{L^2(\Gamma_*)}=\frac{(z_{n,j}-c_{n,j}s_{n,j})+(s_{n,j}-z_{n,j}c_{n,j})c_{n,j}}{z_{n,j}s_{n,j}^2}\|\psi_n(\vartheta,\cdot)\|^2_{\ell^2(V_*)}=\|\psi_n(\vartheta,\cdot)\|^2_{\ell^2(V_*)}.
$$

The identity (1.25) follows from Proposition 3.3. ■

The first eigenvalue $\lambda_1(\vartheta)$ and the corresponding normalized eigenfunction $\psi_1(\vartheta,\cdot)$ have asymptotics as $\vartheta=\varepsilon\omega$, $\varepsilon=|\vartheta\,|\to 0$, $\omega\in\mathbb{S}^{d-1}$:

$$
\begin{aligned}
&\lambda_1(\vartheta)=\lambda_1(0)+\varepsilon^2\mu(\omega)+O(\varepsilon^3),\qquad \lambda_1(0)=-1,\qquad \mu(\omega)=\frac12\,\ddot{\lambda}_1(\varepsilon\omega)\big|_{\varepsilon=0},\\
&\psi_1(\vartheta,\cdot)=\psi_1(0,\cdot)+\varepsilon\psi_1^{(1)}+O(\varepsilon^2),\qquad \psi_1(0,v)=\sqrt{\frac{\varkappa_v}{\varkappa}}\,,\qquad \varkappa=\sum_{v\in V_*}\varkappa_v,\\
&\qquad\qquad\qquad\qquad\qquad\qquad \psi_1^{(1)}=\psi_1^{(1)}(\omega,\cdot)=\dot{\psi}_1(\varepsilon\omega,\cdot)\big|_{\varepsilon=0},
\end{aligned}
\tag{3.34}
$$

where $\dot{g}=\frac{\partial g}{\partial\varepsilon}$ and $\mathbb{S}^d$ is the d-dimensional sphere.

Proposition 3.3. *i) Each eigenfunction* $\Psi_{1,j}(\vartheta)=\big(\Psi_{1,j,\mathbf{e}}(\vartheta,t)\big)_{\mathbf{e}\in\mathcal{E}_*}$*,* $j\in\mathbb{Z}_0$*, of the operator* $\Delta_M(\vartheta)$*,* $\vartheta\in\mathbb{T}^d\setminus\{0\}$*, defined by (1.23), satisfies as* $\mathbf{e}=(u,v)$ *and* $\varepsilon=|\vartheta|\to 0$

$$
\Psi_{1,j,\mathbf{e}}(\vartheta,t)=\frac{(-1)^j\sqrt2}{\sqrt{\varkappa}}\,\cos(\pi j_*t)-\frac{\sin(\pi j_*t)}{\sqrt{\mu(\omega)}}\bigg(\frac{\psi_1^{(1)}(\omega,u)}{\sqrt{\varkappa_u}}-\frac{\psi_1^{(1)}(\omega,v)}{\sqrt{\varkappa_v}}-\frac{i}{\sqrt{\varkappa}}\langle\tau(\mathbf{e}),\omega\rangle\bigg)+O(\varepsilon),
\tag{3.35}
$$

where $j_*=j$*, if* j *is even, and* $j_*=j+1$ *if* j *is odd.*

ii) Let $\vartheta\in\mathbb{T}^d\setminus\{0\}$ *and* $z_1(\vartheta)\leqslant\frac{\pi}{2}$*. Then all eigenfunctions* $\Psi_{1,j}(\vartheta)$*,* $j\in\mathbb{Z}_0$*, of* $\Delta_M(\vartheta)$ *satisfy*

$$
\|\Psi_{1,j}(\vartheta)\|_{L^\infty(\Gamma_*)}<\sqrt2\left(2+\frac{\pi}{\Lambda}+M\,\frac{|\vartheta|}{\sin z_1(\vartheta)}\right),
\tag{3.36}
$$

where

$$
M=\max_{\mathbf{e}\in\mathcal{B}_*}|\tau(\mathbf{e})|+\frac{2}{\Lambda}\max_{u\in V_*}\sum_{\mathbf{e}=(u,v)\in\mathcal{A}_*}\frac{|\tau(\mathbf{e})|}{\sqrt{\varkappa_u\varkappa_v}}\,,
\tag{3.37}
$$

Λ *is the distance between* $\lambda_1(0)=-1$ *and the set* $\sigma\big(\Delta(0)\big)\setminus\big\{\lambda_1(0)\big\}$*.*

iii) Let $\vartheta\in\mathbb{T}^d\setminus\{0\}$ and $z_1(\vartheta)>\frac{\pi}{2}$. Then all eigenfunctions $\Psi_{1,j}(\vartheta)$, $j\in\mathbb{Z}_0$, of $\Delta_M(\vartheta)$ satisfy

$$\|\Psi_{1,j}(\vartheta)\|_{L^\infty(\Gamma_*)}<\frac{2\sqrt{2}}{\sin z_1^+}<\infty, \tag{3.38}$$

where z_1^+ is defined in (1.26).

iv) There exists a constant $C_0>0$ such that

$$\sup_{\vartheta\in\mathbb{T}^d\setminus\{0\}}\frac{|\vartheta|}{\sin z_1(\vartheta)}=C_0<\infty. \tag{3.39}$$

Proof. Let for the shortness

$$z_{1,j}=z_{1,j}(\vartheta),\qquad z_1=z_1(\vartheta),\qquad f_1(u)=\frac{\psi_1(\vartheta,u)}{\sqrt{\varkappa_u}}.$$

i) For the eigenfunction $\Psi_{1,j}(\vartheta)=\big(\Psi_{1,j,\mathbf{e}}(\vartheta,t)\big)_{\mathbf{e}\in\mathcal{E}_*}$, defined by (1.23), we have

$$\begin{aligned}\Psi_{1,j,\mathbf{e}}(\vartheta,t)=\frac{\sqrt{2}}{\sin z_1}\Big[f_1(u)\big(\sin z_{1,j}\cos(z_{1,j}t)-\cos z_{1,j}\sin(z_{1,j}t)\big)\\ +f_1(v)\,e^{i\langle\tau(\mathbf{e}),\,\vartheta\rangle}\sin(z_{1,j}t)\Big]=(-1)^j\sqrt{2}\,f_1(u)\cos(z_{1,j}t)-\sqrt{2}\sin(z_{1,j}t)A(\vartheta),\end{aligned} \tag{3.40}$$

where

$$A(\vartheta)=\frac{1}{\sin z_1}\Big(f_1(u)\cos z_1-f_1(v)\,e^{i\langle\tau(\mathbf{e}),\,\vartheta\rangle}\Big). \tag{3.41}$$

Using (1.22) and (3.34), we have the following asymptotics

$$\begin{aligned}\cos z_1=-\lambda_1(\vartheta)=1-\varepsilon^2\mu(\omega)+O(\varepsilon^3),\\ \sin z_1=\big(1-\lambda_1^2(\vartheta)\big)^{1/2}=\big(2\varepsilon^2\mu(\omega)+O(\varepsilon^3)\big)^{1/2}=\varepsilon\sqrt{2\mu(\omega)}\,\big(1+O(\varepsilon)\big).\end{aligned} \tag{3.42}$$

Substituting (3.34) and (3.42) into (3.41), we obtain

$$\begin{aligned}A(\vartheta)=\frac{(1+O(\varepsilon))}{\varepsilon\sqrt{2\mu(\omega)}}\Big[\frac{\psi_1(0,u)}{\sqrt{\varkappa_u}}+\varepsilon\,\frac{\psi_1^{(1)}(\omega,u)}{\sqrt{\varkappa_u}}-\frac{\psi_1(0,v)}{\sqrt{\varkappa_v}}(1+\varepsilon\, i\langle\tau(\mathbf{e}),\omega\rangle)\\ -\varepsilon\,\frac{\psi_1^{(1)}(\omega,v)}{\sqrt{\varkappa_v}}+O(\varepsilon^2)\Big]=\frac{1}{\sqrt{2\mu(\omega)}}\Big(\frac{\psi_1^{(1)}(\omega,u)}{\sqrt{\varkappa_u}}-\frac{\psi_1^{(1)}(\omega,v)}{\sqrt{\varkappa_v}}-\frac{i}{\sqrt{\varkappa}}\langle\tau(\mathbf{e}),\omega\rangle\Big)+O(\varepsilon).\end{aligned} \tag{3.43}$$

Substituting (3.43) into (3.40) and using (3.34) and

$$\sin(z_{1,j}\,t)=\sin(\pi j_*t)+O(\varepsilon),\qquad \cos(z_{1,j}\,t)=\cos(\pi j_*t)+O(\varepsilon),$$

we get (3.35).

ii) The identity (3.40) gives

$$|\Psi_{1,j,\mathbf{e}}(\vartheta,t)|<\sqrt{2}\big(1+|A(\vartheta)|\big). \tag{3.44}$$

We have the decomposition $\psi_1(\vartheta,v)=\psi_1(0,v)+\widetilde{\psi}_1(\vartheta,v)$ and an identity

$$f_1(u)=f_1^0(u)+\widetilde{f}_1(u),\qquad f_1^0(u)=\frac{\psi_1(0,u)}{\sqrt{\varkappa_u}}=\frac{1}{\sqrt{\varkappa}}. \tag{3.45}$$

We rewrite the difference on the right-hand side of (3.41) in the form

$$\begin{aligned} f_1(u)\cos z_1 - f_1(v)\,e^{i\langle\tau(\mathbf{e}),\,\vartheta\rangle} &= f_1(u) - f_1(v) + f_1(u)(\cos z_1 - 1) - f_1(v)(e^{i\langle\tau(\mathbf{e}),\,\vartheta\rangle} - 1)\\ &= f_1^0(u) - f_1^0(v) + \widetilde f_1(u) - \widetilde f_1(v) - 2f_1(u)\sin^2\frac{z_1}{2} - f_1(v)(e^{i\langle\tau(\mathbf{e}),\,\vartheta\rangle} - 1), \end{aligned}$$

which yields

$$\begin{aligned} \Big|f_1(u)\cos z_1 - f_1(v)\,e^{i\langle\tau(\mathbf{e}),\,\vartheta\rangle}\Big| &< |\widetilde f_1(u)| + |\widetilde f_1(v)| + 2\sin^2\frac{z_1}{2} + 2\Big|\sin\frac{\langle\tau(\mathbf{e}),\,\vartheta\rangle}{2}\Big|\\ &\leqslant |\widetilde f_1(u)| + |\widetilde f_1(v)| + 2\sin^2\frac{z_1}{2} + \big|\langle\tau(\mathbf{e}),\,\vartheta\rangle\big|. \end{aligned} \tag{3.46}$$

Now we estimate $\widetilde\psi_1(\vartheta,\cdot)$, which is defined by the decompositions

$$\Delta(\vartheta) = \Delta(0) + \widetilde\Delta(\vartheta), \qquad \lambda_1(\vartheta) = \lambda_1(0) + \widetilde\lambda_1(\vartheta), \qquad \psi_1(\vartheta,\cdot) = \psi_1(0,\cdot) + \widetilde\psi_1(\vartheta,\cdot).$$

Let P be the orthogonal projection onto the subspace of $\ell^2(V_*)$ orthogonal to $\psi_1(0,\cdot)$. Then we can rewrite the equation $\Delta(\vartheta)\psi_1(\vartheta,\cdot) = \lambda_1(\vartheta)\psi_1(\vartheta,\cdot)$ in the form

$$\big(\Delta(0) + \widetilde\Delta(\vartheta)\big)\big(\psi_1(0,\cdot) + \widetilde\psi_1(\vartheta,\cdot)\big) = \big(\lambda_1(0) + \widetilde\lambda_1(\vartheta)\big)\big(\psi_1(0,\cdot) + \widetilde\psi_1(\vartheta,\cdot)\big) \tag{3.47}$$

or, since $\psi_1(0,\cdot)$ and $\widetilde\psi_1(\vartheta,\cdot)$ are orthogonal,

$$G\widetilde\psi_1(\vartheta,\cdot) = \big(\widetilde\Delta(\vartheta) - \widetilde\lambda_1(\vartheta)1\!\!1_\nu\big)\psi_1(\vartheta,\cdot), \quad \text{where} \quad G = P\big(\Delta(0) - \lambda_1(0)1\!\!1_\nu\big), \tag{3.48}$$

and $1\!\!1_\nu$ is the identity $\nu\times\nu$ matrix. From (3.48) we obtain

$$\widetilde\psi_1(\vartheta,\cdot) = G^{-1}P\big(\widetilde\Delta(\vartheta) - \widetilde\lambda_1(\vartheta)1\!\!1_\nu\big)\psi_1(\vartheta,\cdot). \tag{3.49}$$

This yields

$$\|\widetilde\psi_1(\vartheta,\cdot)\| \leqslant \big\|G^{-1}P\big\|\cdot\big\|\big(\widetilde\Delta(\vartheta) - \widetilde\lambda_1(\vartheta)1\!\!1_\nu\big)\psi_1(\vartheta,\cdot)\big\| \leqslant \frac{1}{\Lambda}\,\|\widetilde\Delta(\vartheta) - \widetilde\lambda_1(\vartheta)1\!\!1_\nu\|, \tag{3.50}$$

where Λ is the distance between $\lambda_1(0) = -1$ and $\sigma\big(\Delta(0)\big)\setminus\big\{\lambda_1(0)\big\}$. Due to (2.5), we have

$$\|\widetilde\Delta(\vartheta) - \widetilde\lambda_1(\vartheta)1\!\!1_\nu\| \leqslant \widetilde\lambda_1(\vartheta) + \max_{u\in V_*}\sum_{\mathbf{e}=(u,v)\in\mathcal{A}_*}\frac{\big|\langle\tau(\mathbf{e}),\,\vartheta\rangle\big|}{\sqrt{\varkappa_u\varkappa_v}} \leqslant \widetilde\lambda_1(\vartheta) + |\vartheta|\,T, \tag{3.51}$$

where $T = \max\limits_{u\in V_*}\sum\limits_{\mathbf{e}=(u,v)\in\mathcal{A}_*}\frac{|\tau(\mathbf{e})|}{\sqrt{\varkappa_u\varkappa_v}}$. Substituting (3.51) into (3.50), we obtain

$$\|\widetilde\psi_1(\vartheta,\cdot)\| \leqslant \frac{1}{\Lambda}\big(\widetilde\lambda_1(\vartheta) + |\vartheta|\,T\big). \tag{3.52}$$

Using (3.46), (3.52), for $A(\vartheta)$ defined by (3.41), we have

$$\begin{aligned} |A(\vartheta)| &\leqslant \frac{1}{\sin z_1}\bigg(\frac{2}{\Lambda}\Big(\widetilde\lambda_1(\vartheta) + |\vartheta|\,T\Big) + 2\sin^2\frac{z_1}{2} + \big|\langle\tau(\mathbf{e}),\,\vartheta\rangle\big|\bigg)\\ &= \tan\frac{z_1}{2} + \frac{2}{\Lambda}\,\frac{\widetilde\lambda_1(\vartheta)}{\sin z_1} + \frac{|\vartheta|}{\sin z_1}\bigg(\frac{2}{\Lambda}\,T + \max_{\mathbf{e}\in\mathcal{B}_*}|\tau(\mathbf{e})|\bigg). \end{aligned} \tag{3.53}$$

Let $z_1\in(0,\frac{\pi}{2}]$. Then $\sin z_1 \geqslant \frac{2}{\pi}\,z_1$. This and (3.53) give

$$|A(\vartheta)| \leqslant 1 + \frac{\pi}{\Lambda}\cdot\frac{\widetilde\lambda_1(\vartheta)}{z_1(\vartheta)} + M\,\frac{|\vartheta|}{\sin z_1(\vartheta)}\,, \tag{3.54}$$

where M is defined by (3.37). Due to $z_1(0)=0$ we have the simple inequality

$$\widetilde{\lambda}_1(\vartheta)=\lambda_1(\vartheta)-\lambda_1(0)=-\cos z_1(\vartheta)+1=\int\limits_0^{z_1(\vartheta)}\sin t\,dt\leqslant z_1(\vartheta),$$

which yields that $\dfrac{\widetilde{\lambda}_1(\vartheta)}{z_1(\vartheta)}\leqslant 1$. Then the estimate (3.54) has the form

$$|A(\vartheta)|\leqslant 1+\frac{\pi}{\Lambda}+M\,\frac{|\vartheta|}{\sin z_1(\vartheta)}\,. \tag{3.55}$$

Combining (3.44) and (3.55), we obtain (3.36).

iii) We have $\frac{\pi}{2}<z_1(\vartheta)\leqslant z_1^+<\pi$. This yields $0<\sin z_1^+<\sin z_1(\vartheta)$. We note that $z_1^+\neq\pi$, since a bipartite graph Γ_* consists of more than 1 vertex. Then the estimate (3.38) follows directly from the identity (1.23).

iv) Using the second formula in (3.42), we obtain

$$\frac{|\vartheta|}{\sin z_1(\vartheta)}=\frac{1}{\sqrt{2\mu(\omega)}}+O\big(|\vartheta|\big)\qquad\text{as}\quad|\vartheta|\to 0. \tag{3.56}$$

The following estimate holds true (see Theorem 1.2 in [KS15c]):

$$\begin{aligned}&\mu(\omega)\geqslant T_1=\frac{1}{\varkappa\,\nu\,d}\left(\frac{d-1}{C}\right)^{d-1}>0,\qquad\forall\,\omega\in\mathbb{S}^{d-1},\\&\varkappa=\sum_{v\in V_*}\varkappa_v,\qquad C=\sum_{j=1}^d|\tau(\mathbf{e}_j)|^2,\end{aligned} \tag{3.57}$$

where $\mathbf{e}_1,\ldots,\mathbf{e}_d\in\mathcal{B}_*$ are bridges with linearly independent indices $\tau(\mathbf{e}_1),\ldots,\tau(\mathbf{e}_d)\in\mathbb{Z}^d$. Substituting the estimate (3.57) into (3.56), we get

$$\frac{|\vartheta|}{\sin z_1(\vartheta)}\leqslant\frac{1}{\sqrt{2T_1}}+O\big(|\vartheta|\big).$$

Then there exists a constant $C_1>0$ such that

$$\frac{|\vartheta|}{\sin z_1(\vartheta)}<\frac{1}{\sqrt{2T_1}}+C_1,\quad\text{for all}\quad|\vartheta|<R, \tag{3.58}$$

where R is some positive number. Now let $|\vartheta|\geqslant R$. Then $\sin z_1(\vartheta)\geqslant\alpha$ for some constant $\alpha>0$ and

$$\frac{|\vartheta|}{\sin z_1(\vartheta)}\leqslant\frac{\pi\sqrt{d}}{\alpha}<\infty. \tag{3.59}$$

Combining (3.58) and (3.59), we obtain (3.39). ■

Proposition 3.4. *i) All eigenfunctions* $\Psi_{n,j}(\vartheta)$, $n=2,\ldots,\nu-1$, $(j,\vartheta)\in\mathbb{Z}_0\times(\mathbb{T}^d\setminus\{0\})$, *satisfy*

$$\|\Psi_{n,j}(\vartheta)\|_{L^\infty(\Gamma_*)}<\frac{2\sqrt{2}}{\alpha}\,,\qquad\alpha=\inf_{\substack{n=2,\ldots,\nu-1\\ \vartheta\in\mathbb{T}^d}}\sin z_n(\vartheta)>0. \tag{3.60}$$

ii) Let $\vartheta\in\mathbb{T}^d\setminus\{0\}$ and let Γ_* be non-bipartite. Then all eigenfunctions $\Psi_{\nu,j}(\vartheta)$, $j\in\mathbb{Z}_0$, of $\Delta_M(\vartheta)$ satisfy

$$\|\Psi_{\nu,j}(\vartheta)\|_{L^\infty(\Gamma_*)}<\frac{2\sqrt{2}}{\alpha_\nu},\qquad \alpha_\nu=\inf_{\vartheta\in\mathbb{T}^d}\sin z_\nu(\vartheta)>0. \tag{3.61}$$

iii) Let $\vartheta\in\mathbb{T}^d\setminus\{0\}$ and let Γ_* be bipartite. Then all eigenfunctions $\Psi_{\nu,j}(\vartheta)$, $j\in\mathbb{Z}_0$, of $\Delta_M(\vartheta)$ satisfy

$$\|\Psi_{\nu,j}(\vartheta)\|_{L^\infty(\Gamma_*)}<\sqrt{2}\left(2+\frac{\pi}{\Lambda}+M\,\frac{|\vartheta|}{\sin z_1(\vartheta)}\right), \tag{3.62}$$

where M and Λ are defined in Proposition 3.3.ii.

Proof. i) This estimate follows from the identity (1.23).

ii) Since Γ is non-bipartite, using (1.23), we also have (3.61).

iii) Let for the shortness

$$z_{\nu,j}=z_{\nu,j}(\vartheta),\qquad z_\nu=z_\nu(\vartheta),\qquad f_\nu(u)=\frac{\psi_\nu(\vartheta,u)}{\sqrt{\varkappa_u}}.$$

For the eigenfunction $\Psi_{\nu,j}(\vartheta)=\big(\Psi_{\nu,j,\mathbf{e}}(\vartheta,t)\big)_{\mathbf{e}\in\mathcal{E}_*}$, defined by (1.23), we have

$$\Psi_{\nu,j,\mathbf{e}}(\vartheta,t)=(-1)^j\sqrt{2}\,f_\nu(u)\cos(z_{\nu,j}t)-\sqrt{2}\sin(z_{\nu,j}t)A_\nu(\vartheta), \tag{3.63}$$

where

$$A_\nu(\vartheta)=\frac{1}{\sin z_\nu}\Big(f_\nu(u)\cos z_\nu-f_\nu(v)\,e^{i\langle\tau(\mathbf{e}),\vartheta\rangle}\Big).$$

For bipartite graph Γ_* with parts V_1 and V_2 we have

$$\sin z_\nu=\sin z_1,\quad \cos z_\nu=-\cos z_1,\qquad f_\nu(v)=\begin{cases}f_1(v), & \text{if } v\in V_1\\ -f_1(v), & \text{if } v\in V_2\end{cases}.$$

This yields that $A_\nu(\vartheta)=A(\vartheta)$, where $A(\vartheta)$ is defined by (3.41). Combining this, (3.63) and (3.55), we obtain (3.62). ∎

Remark. The estimates (3.36), (3.38), (3.60) – (3.62) give that the eigenfunctions $\Psi_{n,j}(\vartheta)$, $(n,j,\vartheta)\in\mathbb{N}_\nu\times\mathbb{Z}_0\times\mathbb{T}^d\setminus\{0\}$, are uniformly bounded with respect to n,j,ϑ.

Proof of Corollary 1.4. Recall that $\Delta_M=\Delta_{MD}\oplus\Delta_{MV}$ and Theorem 1.2 gives

$$\sigma(\Delta_{MD})=\sigma_{fb}(\Delta_{MD})=\sigma_D. \tag{3.64}$$

Thus, we need to consider the operator Δ_{MV}. Since each $\lambda_n(\cdot)$ is a real and piecewise analytic function on the torus $\mathbb{T}^d$, each $z_{n,j}(\cdot)$, $(n,j)\in\mathbb{N}_\nu\times\mathbb{Z}_0$, defined by the formula (1.22), is also a real and piecewise analytic function on $\mathbb{T}^d$. Then (1.22) yields

$$\sigma(\Delta_{MV})=\bigcup_{(n,j)\in\mathbb{N}_\nu\times\mathbb{Z}_0}\sigma_{n,j}(\Delta_M),\qquad \sigma_{n,j}(\Delta_M)=[E_{n,j}^-,E_{n,j}^+]=z_{n,j}^2(\mathbb{T}^d), \tag{3.65}$$

where $E_{n,j}^\pm$ are defined in (1.26). Since $\lambda_n(\cdot)=\text{const}$, for $\nu-r<n\leqslant\nu$ and all other eigenvalues $\lambda_1(\cdot),\ldots,\lambda_{\nu-r}(\cdot)$ are not constants (in any small ball), we obtain (1.26). ∎

Recall that we identify an edge $\mathbf{e}_*=(u_*,v_*)\in\mathcal{E}_*$ of the fundamental graph $\Gamma_*=(V_*,\mathcal{E}_*)$ with an index $\tau(\mathbf{e}_*)$ with the edge $\mathbf{e}=(u,v+\tau(\mathbf{e}_*))\in\mathcal{E}$ of the graph $\Gamma=(V,\mathcal{E})$, where $u,v\in V_0$ such that $u_*=\mathfrak{f}_V(u)$, $v_*=\mathfrak{f}_V(v)$.

Proof of Proposition 1.5. i) We fix $\vartheta\in\mathbb{T}^d\setminus\{0\}$ and consider the space X of all functions $f:V\to\mathbb{C}$ satisfying the following quasiperiodic condition:

$$f(v+m)=e^{i\langle m,\,\vartheta\rangle}f(v),\qquad\text{for all}\quad (v,m)\in V\times\mathbb{Z}^d. \tag{3.66}$$

Such a function f is uniquely determined by its values at the fundamental graph vertices. Thus, X is isomorphic to $\ell^2(V_*)$. The fiber Laplacian $\Delta(\vartheta):\ell^2(V_*)\to\ell^2(V_*)$ can be also defined as the restriction to the space X of the operator Δ.

Let $\psi_n(\vartheta,\cdot)\in\ell^2(V_*)$ be the normalized eigenfunction of $\Delta(\vartheta)$ corresponding to the eigenvalue $\lambda_n(\vartheta)$. The function $\widetilde\psi_n(\vartheta,\cdot)$ defined by (1.28) satisfies the quasiperiodic condition:

$$\widetilde\psi_n(\vartheta,v+m)=e^{i\langle m,\,\vartheta\rangle}\widetilde\psi_n(\vartheta,v),\qquad (v,m)\in V\times\mathbb{Z}^d, \tag{3.67}$$

i.e., $\widetilde\psi_n(\vartheta,\cdot)\in X$. For each $v\in V$ we have the unique representation $v=v_0+m$ for some $v_0\in V_0$ and $m\in\mathbb{Z}^d$ and, due to (3.67) and the definition of edge indices, we obtain

$$\begin{aligned}
&(\Delta\widetilde\psi_n(\vartheta,\cdot))(v)=(\Delta(\vartheta)\widetilde\psi_n(\vartheta,\cdot))(v)=e^{i\langle m,\,\vartheta\rangle}(\Delta(\vartheta)\widetilde\psi_n(\vartheta,\cdot))(v_0)\\
&=-\frac{e^{i\langle m,\,\vartheta\rangle}}{\sqrt{\varkappa_{v_0}}}\sum_{(v_0,\,u)_e\in\mathcal{E}}\frac{1}{\sqrt{\varkappa_u}}\widetilde\psi_n(\vartheta,u)=-\frac{e^{i\langle m,\,\vartheta\rangle}}{\sqrt{\varkappa_{v_0}}}\sum_{\mathbf{e}=(v_0,\,u_0)_e\in\mathcal{E}_*}\frac{e^{i\langle\tau(\mathbf{e}),\,\vartheta\rangle}}{\sqrt{\varkappa_{u_0}}}\,\widetilde\psi_n(\vartheta,u_0)\\
&=-\frac{e^{i\langle m,\,\vartheta\rangle}}{\sqrt{\varkappa_{v_0}}}\sum_{\mathbf{e}=(v_0,\,u_0)_e\in\mathcal{E}_*}\frac{e^{i\langle\tau(\mathbf{e}),\,\vartheta\rangle}}{\sqrt{\varkappa_{u_0}}}\,\psi_n(\vartheta,u_0)=e^{i\langle m,\,\vartheta\rangle}(\Delta(\vartheta)\psi_n(\vartheta,\cdot))(v_0)\\
&=e^{i\langle m,\,\vartheta\rangle}\lambda_n(\vartheta)\psi_n(\vartheta,v_0)=\lambda_n(\vartheta)\widetilde\psi_n(\vartheta,v).
\end{aligned}$$

Thus, $\widetilde\psi_n(\vartheta,\cdot)\in\ell^2(V)$ is an eigenfunction of Δ corresponding to the eigenvalue $\lambda_n(\vartheta)$.

ii) The proof of this item is similar to the proof of the item i).

iii) If we denote $u=v+m$ in (3.67), where $u\in V\subset\mathbb{R}^d$, then for all $v\in V$ (1.28) gives

$$\widetilde\psi_n(\vartheta,u)=e^{i\langle u-v,\,\vartheta\rangle}\widetilde\psi_n(\vartheta,u-m)=e^{i\langle u,\,\vartheta\rangle}e^{-i\langle v+m,\,\vartheta\rangle}\widetilde\psi_n(\vartheta,u)=e^{i\langle u,\,\vartheta\rangle}\varphi_n(\vartheta,u).$$

The function $\varphi_n(\vartheta,u)=e^{-i\langle u,\,\vartheta\rangle}\widetilde\psi_n(\vartheta,u)$ is $\mathbb{Z}^d$-periodic with respect to u, since

$$\begin{aligned}
\varphi_n(\vartheta,u+m)&=e^{-i\langle u+m,\,\vartheta\rangle}\widetilde\psi_n(\vartheta,u+m)=e^{-i\langle u+m,\,\vartheta\rangle}e^{i\langle m,\,\vartheta\rangle}\widetilde\psi_n(\vartheta,u)\\
&=e^{-i\langle u,\,\vartheta\rangle}\widetilde\psi_n(\vartheta,u)=\varphi_n(\vartheta,u),\qquad (u,m)\in V\times\mathbb{Z}^d.
\end{aligned}$$

Thus, (1.30) has been proved. The proof of the identity (1.31) is similar. ■

We recall known results (see Theorems 4.5.2 and 4.5.4 in [BK13]) about the connection between true eigenfunctions from $\ell^2(V)$ and $L^2(\Gamma)$ of the discrete and metric Laplacians, respectively, and eigenfunctions of their fiber operators.

Proposition 3.5. *Let $\psi_n(\vartheta,\cdot)\in\ell^2(V_*)$, $\vartheta\in\mathbb{T}^d\setminus\{0\}$, be the family of the normalized eigenfunctions of the fiber operators $\Delta(\vartheta)$ corresponding to the eigenvalue $\lambda_n\in\sigma_{fb}(\Delta)$ for some $n=\nu-r+1,\ldots,\nu$ and let $\Psi_{n,j}(\vartheta,\cdot)\in L^2(\Gamma_*)$ be the family of the normalized eigenfunctions of the fiber operators $\Delta_M(\vartheta)$ corresponding to the eigenvalue $z_{n,j}^2\in\sigma_{fb}(\Delta_M)$ defined by (1.22), $j=0,1,2,\ldots$. Then the following statements hold true.*

i) The eigenfunction $\widetilde{\psi}_n(\cdot)\in\ell^2(V)$ *of the discrete Laplacian* Δ *corresponding to the flat band* λ_n *has the form*

$$\widetilde{\psi}_n(v+m)=\int_{\mathbb{T}^d}e^{i\langle m,\vartheta\rangle}\psi_n(\vartheta,v)\frac{d\vartheta}{(2\pi)^d}\,,\qquad\forall\,(v,m)\in V_*\times\mathbb{Z}^d. \tag{3.68}$$

ii) The eigenfunction $\widetilde{\Psi}_{n,j}(\cdot)\in L^2(\Gamma)$ *of the metric Laplacian* Δ_M *corresponding to the flat band* $z_{n,j}^2$ *has the form*

$$\widetilde{\Psi}_{n,j}(x+m)=\int_{\mathbb{T}^d}e^{i\langle m,\vartheta\rangle}\Psi_{n,j}(\vartheta,x)\frac{d\vartheta}{(2\pi)^d}\,,\qquad\forall\,(x,m)\in\Gamma_*\times\mathbb{Z}^d. \tag{3.69}$$

Remarks. 1) The right-hand sides of (3.68) and (3.69) are the inverse operators of the unitary operators (Floquet transforms) U and $\mathscr{U}$ from (1.4) and (1.11), respectively.

2) The eigenfunctions defined by (3.68) and (3.69) are compactly supported.

4. SCHRÖDINGER OPERATORS ON PERIODIC METRIC GRAPHS

4.1. **Trace class estimates.** We consider Schrödinger operators H on the periodic graph Γ given by

$$H=H_0+Q,\qquad H_0=\Delta_M.$$

We assume that the potential $Q=(Q_{\mathbf{e}})_{\mathbf{e}\in\mathcal{E}}$ is real and belongs to the spaces $L^1(\Gamma)\cap L^2(\Gamma)$. Recall that $R_0(k)=(H_0-k^2)^{-1}$, $R(k)=(H-k^2)^{-1}$ and

$$Y_0(k)=|Q|^{\frac12}\,R_0(k)\,Q^{1/2},\qquad Y(k)=|Q|^{\frac12}\,R(k)\,Q^{\frac12},\quad Q^{\frac12}=|Q|^{\frac12}\operatorname{sign}Q, \tag{4.1}$$

$k\in\mathbb{C}_+$. Note that Propositions 3.3 and 3.4 give

$$C_\Gamma:=\sup_{(n,j,\vartheta)\in\mathbb{N}_\nu\times\mathbb{Z}_0\times\mathbb{T}^d}\|\Psi_{n,j}(\vartheta)\|_{L^\infty(\Gamma_*)}<\infty. \tag{4.2}$$

Let $\mathcal{R}_k(x,x',\vartheta),x,x'\in\Gamma_*$, be the kernel of the integral operator $(\Delta_{MV}(\vartheta)-k^2)^{-1}$ on $L^2(\Gamma_*)$. Due to Theorem 1.3 we have

$$\mathcal{R}_k(x,x',\vartheta)=\sum_{j=0}^{\infty}\sum_{n=1}^{\nu}\frac{\Psi_{n,j}(\vartheta,x)\overline{\Psi}_{n,j}(\vartheta,x')}{z_{n,j}^2(\vartheta)-k^2}\,,\qquad x,x'\in\Gamma_*. \tag{4.3}$$

Theorem 4.1. *i) Let* $(Q,k)\in L^1(\Gamma)\times\mathbb{C}_+$ *and let* $\Delta_M^{\frac12}\geqslant 0$. *Then* $\big(\Delta_M^{\frac12}-k\big)^{-1}|Q|^{\frac12}\in\mathbf{B}_2$ *and* $Y_0(k)\in\mathbf{B}_1$ *and they satisfy*

$$\|(\Delta_M^{\frac12}-k)^{-1}|Q|^{\frac12}\|_{\mathbf{B}_2}^2\leqslant C_0\|Q\|_{L^1(\Gamma)}\bigg(\frac{2}{(\operatorname{Im}k)^2}+\frac{1}{\operatorname{Im}k}\bigg), \tag{4.4}$$

$$\|Y_0(k)\|_{\mathbf{B}_1}\leqslant\mathcal{C}(Q,k),$$

$$\text{where } \mathcal{C}(Q,k)=C_0\|Q\|_{L^1(\Gamma)}\bigg(\frac{2}{(\operatorname{Im}k)^2}+\frac{1}{\operatorname{Im}k}\bigg)^{\frac12}\bigg(\frac{2}{|k|^2}+\frac{1}{|k|}\bigg)^{\frac12}, \tag{4.5}$$

$$C_0=2(\nu C_\Gamma^2+\nu_*-\nu),$$

$$\|Y_0(k)\|_{\mathbf{B}_1}=\frac{O(1)}{|k\operatorname{Im}k|^{\frac12}}\qquad\textit{as}\qquad\operatorname{Im}k\to\infty. \tag{4.6}$$

Moreover, if $\operatorname{Re} k \leqslant 0$, *then*

$$\|(\Delta_M^{\frac12}-k)^{-1}|Q|^{\frac12}\|_{\mathbf{B}_2}^2 \leqslant C_0\|Q\|_{L^1(\Gamma)}\biggl(\frac{2}{|k|^2}+\frac{1}{|k|}\biggr). \tag{4.7}$$

ii) Let, in addition,

$$\sup_{x\in\Gamma_*}\bigl|Q(x+m)\bigr| = \gamma_m, \quad m\in\mathbb{Z}^d, \qquad \sum_{m\in\mathbb{Z}^d}\gamma_m = \gamma < \infty. \tag{4.8}$$

Then

$$\|(\Delta_M^{\frac12}-k)^{-1}|Q|^{\frac12}\|_{\mathbf{B}_2}^2 \leqslant \gamma(\nu_*+\nu)\biggl(\frac{2}{(\operatorname{Im} k)^2}+\frac{1}{\operatorname{Im} k}\biggr). \tag{4.9}$$

Moreover, if $\operatorname{Re} k \leqslant 0$, *then*

$$\|(\Delta_M^{\frac12}-k)^{-1}|Q|^{\frac12}\|_{\mathbf{B}_2}^2 \leqslant \gamma(\nu_*+\nu)\biggl(\frac{2}{|k|^2}+\frac{1}{|k|}\biggr). \tag{4.10}$$

Proof. i) Due to Theorem 1.1, the operator $(\Delta_M^{\frac12}-k)^{-1}$ has the form given by

$$\Bigl(\mathscr{U}(\Delta_M^{\frac12}-k)^{-1}h\Bigr)(\vartheta,\cdot) = \Bigl(\bigl(\Delta_{MD}^{\frac12}(\vartheta)-k\bigr)^{-1}\oplus\bigl(\Delta_{MV}^{\frac12}(\vartheta)-k\bigr)^{-1}\Bigr)(\mathscr{U}h)(\vartheta,\cdot), \tag{4.11}$$

where the unitary operator $\mathscr{U}: L^2(\Gamma)\to\mathscr{H}$ is defined by (1.11), $h\in L^2(\Gamma)$ is compactly supported and $\vartheta\in\mathbb{T}^d$. Consider the second term on the right-hand side of (4.11). Using Theorem 1.3 and denoting $\alpha=(n,j)\in\mathbb{N}_\nu\times\mathbb{Z}_0$, $\mathbb{Z}_0=\{0,1,2,\ldots\}$, we obtain

$$\begin{aligned}\bigl(\Delta_{MV}^{\frac12}(\vartheta)-k\bigr)^{-1}(\mathscr{U}h)(\vartheta,x) &= \sum_{\alpha\in\mathbb{N}_\nu\times\mathbb{Z}_0}\frac{1}{z_\alpha(\vartheta)-k}\,\mathcal{P}_\alpha(\vartheta)(\mathscr{U}h)(\vartheta,x)\\ &= \sum_{\alpha\in\mathbb{N}_\nu\times\mathbb{Z}_0}\frac{1}{z_\alpha(\vartheta)-k}\,\Psi_\alpha(\vartheta,x)\int_{\Gamma_*}(\mathscr{U}h)(\vartheta,x')\overline{\Psi}_\alpha(\vartheta,x')\,dx'.\end{aligned}$$

Using $Q\in L^1(\Gamma)$, the normalized condition $\int_{\Gamma_*}\bigl|\Psi_\alpha(\vartheta,x)\bigr|^2dx=1$ and (4.2), this gives

$$\begin{aligned}&\|\bigl(\Delta_{MV}^{\frac12}(\cdot)-k\bigr)^{-1}\mathscr{U}|Q|^{\frac12}\|_{\mathbf{B}_2}^2\\ &= \frac{1}{(2\pi)^d}\sum_{m\in\mathbb{Z}^d}\sum_{\alpha\in\mathbb{N}_\nu\times\mathbb{Z}_0}\int_{\mathbb{T}^d}\frac{d\vartheta}{|z_\alpha(\vartheta)-k|^2}\int_{\Gamma_*}\bigl|Q(x+m)\bigr|\cdot\bigl|\Psi_\alpha(\vartheta,x)\bigr|^2dx\\ &\leqslant \frac{C_\Gamma^2}{(2\pi)^d}\sum_{m\in\mathbb{Z}^d}\sum_{\alpha\in\mathbb{N}_\nu\times\mathbb{Z}_0}\int_{\mathbb{T}^d}\frac{d\vartheta}{|z_\alpha(\vartheta)-k|^2}\int_{\Gamma_*}\bigl|Q(x'+m)\bigr|\,dx'\\ &= \frac{C_\Gamma^2}{(2\pi)^d}\,\|Q\|_{L^1(\Gamma)}\sum_{\alpha\in\mathbb{N}_\nu\times\mathbb{Z}_0}\int_{\mathbb{T}^d}\frac{d\vartheta}{|z_\alpha(\vartheta)-k|^2}\,.\end{aligned} \tag{4.12}$$

Let $k=p+iq\in\mathbb{C}$, $q\neq0$. We assume that the following inequality holds true

$$\sum_{j=0}^{\infty}\frac{1}{|z_{n,j}(\vartheta)-k|^2}\leqslant\frac{4}{q^2}+\frac{2}{|q|} \qquad \text{for each } \ n\in\mathbb{N}_\nu. \tag{4.13}$$

Substituting (4.13) into (4.12), we obtain

$$\|\bigl(\Delta_{MV}^{\frac12}(\cdot)-k\bigr)^{-1}\mathscr{U}|Q|^{\frac12}\|_{\mathbf{B}_2}^2 \leqslant 2\nu\,C_\Gamma^2\,\|Q\|_{L^1(\Gamma)}\biggl(\frac{2}{q^2}+\frac{1}{|q|}\biggr). \tag{4.14}$$

Now we prove (4.13). Due to (1.22), we have the following identity

$$\sum_{j=0}^{\infty}\frac{1}{|z_{n,j}(\vartheta)-k|^2}=\sum_{\substack{j=0\\ j\text{ is even}}}^{\infty}\frac{1}{\big(z_n(\vartheta)+\pi j-p\big)^2+q^2}+\sum_{\substack{j=1\\ j\text{ is odd}}}^{\infty}\frac{1}{\big(\pi(j+1)-z_n(\vartheta)-p\big)^2+q^2}\,.$$

Let

$$p-z_n(\vartheta)=\pi(m+a),\qquad p+z_n(\vartheta)=\pi(m_1+a_1)$$

for some $m,m_1\in\mathbb{Z}$, and $a=a(\vartheta),a_1=a_1(\vartheta)\in(0,1)$. Then

$$\begin{aligned}\sum_{j=0}^{\infty}\frac{1}{|z_{n,j}(\vartheta)-k|^2}&=\sum_{\substack{j=0\\ j\text{ is even}}}^{\infty}\frac{1}{\pi^2(j-m-a)^2+q^2}+\sum_{\substack{j=1\\ j\text{ is odd}}}^{\infty}\frac{1}{\pi^2(j+1-m_1-a_1)^2+q^2}\\ &\leqslant\sum_{j=-\infty}^{\infty}\frac{1}{\pi^2(j-a)^2+q^2}+\sum_{j=-\infty}^{\infty}\frac{1}{\pi^2(j-a_1)^2+q^2}\leqslant\frac{4}{q^2}+\sum_{j=1}^{\infty}\frac{4}{\pi^2j^2+q^2}\\ &\leqslant\frac{4}{q^2}+4\int_0^{\infty}\frac{dt}{\pi^2t^2+q^2}=\frac{4}{q^2}+\frac{2}{|q|}\,.\end{aligned}\tag{4.15}$$

Consider the first term on the right-hand side of (4.11). Similar arguments yield

$$\|\big(\Delta_{MD}^{\frac12}(\cdot)-k\big)^{-1}\mathscr{U}|Q|^{\frac12}\|_{\mathbf{B}_2}^2\leqslant 2(\nu_*-\nu)\,\|Q\|_{L^1(\Gamma)}\bigg(\frac{2}{q^2}+\frac{1}{|q|}\bigg).\tag{4.16}$$

Summing (4.14) and (4.16) and using (4.11), we get

$$\begin{aligned}\|(\Delta_M^{\frac12}-k)^{-1}|Q|^{\frac12}\|_{\mathbf{B}_2}^2&=\|\mathscr{U}(\Delta_M^{\frac12}-k)^{-1}|Q|^{\frac12}\|_{\mathbf{B}_2}^2\\ \leqslant 2(\nu\,C_\Gamma^2+\nu_*-\nu)\|Q\|_{L^1(\Gamma)}\bigg(\frac{2}{q^2}+\frac{1}{|q|}\bigg)&=C_0\|Q\|_{L^1(\Gamma)}\bigg(\frac{2}{(\operatorname{Im}k)^2}+\frac{1}{|\operatorname{Im}k|}\bigg).\end{aligned}\tag{4.17}$$

Thus, (4.4) has been proved.

Let, in addition, $\operatorname{Re}k\leqslant 0$. Then

$$|z_\alpha(\vartheta)-k|^2=\big(z_\alpha(\vartheta)-p\big)^2+q^2\geqslant z_\alpha^2(\vartheta)+p^2+q^2=z_\alpha^2(\vartheta)+|k|^2\tag{4.18}$$

and using above arguments we obtain (4.7).

For any $k\in\mathbb{C}_+$ we have the following factorization

$$Y_0(k)=|Q|^{\frac12}R_0(k)Q^{\frac12}=\big(|Q|^{\frac12}(\Delta_M^{\frac12}+k)^{-1}\big)\big((\Delta_M^{\frac12}-k)^{-1}Q^{\frac12}\big).$$

Then

$$\|Y_0(k)\|_{\mathbf{B}_1}\leqslant\big\||Q|^{\frac12}(\Delta_M^{\frac12}+k)^{-1}\big\|_{\mathbf{B}_2}\big\|(\Delta_M^{\frac12}-k)^{-1}Q^{\frac12}\big\|_{\mathbf{B}_2}$$

and, applying the estimates (4.17) and (4.7), we obtain (4.5), which yields asymptotics (4.6).

ii) Let the potential Q satisfy (4.8). Then using (4.12) and (4.13), we obtain

$$\begin{aligned}&\|\big(\Delta_{MV}^{\frac12}(\cdot)-k\big)^{-1}\mathscr{U}|Q|^{\frac12}\|_{\mathbf{B}_2}^2\\ &=\frac{1}{(2\pi)^d}\sum_{m\in\mathbb{Z}^d}\sum_{\alpha\in\mathbb{N}_\nu\times\mathbb{Z}_0}\int_{\mathbb{T}^d}\frac{1}{|z_\alpha(\vartheta)-k|^2}\int_{\Gamma_*}\big|Q(x'+m)\big|\cdot\big|\Psi_\alpha(\vartheta,x')\big|^2dx'\,d\vartheta\\ &\leqslant\frac{1}{(2\pi)^d}\sum_{m\in\mathbb{Z}^d}\gamma_m\sum_{\alpha\in\mathbb{N}_\nu\times\mathbb{Z}_0}\int_{\mathbb{T}^d}\frac{d\vartheta}{|z_\alpha(\vartheta)-k|^2}\leqslant 2\gamma\,\nu\bigg(\frac{2}{(\operatorname{Im}k)^2}+\frac{1}{|\operatorname{Im}k|}\bigg).\end{aligned}\tag{4.19}$$

Similar arguments yield

$$\|\big(\Delta_{MD}^{\frac{1}{2}}(\cdot)-k\big)^{-1}\mathscr{U}|Q|^{\frac{1}{2}}\|_{\mathbf{B}_2}^2 \leqslant \gamma(\nu_*-\nu)\bigg(\frac{2}{(\operatorname{Im}k)^2}+\frac{1}{|\operatorname{Im}k|}\bigg). \tag{4.20}$$

Summing (4.19) and (4.20) and using (4.11), we get (4.9).

Let, in addition, $\operatorname{Re}k\leqslant 0$. Then using (4.18) and above arguments we obtain (4.10). ■

Lemma 4.2. *Let* $(Q,k)\in L^1(\Gamma)\times\mathbb{C}_+$. *Then*

$$\operatorname{Tr}Y_0(k)=\frac{1}{(2\pi)^d}\int_{\mathbb{T}^d}\bigg(\sum_{j=0}^{\infty}\sum_{n=1}^{\nu}\frac{Q_{n,j}(\vartheta)}{z_\alpha^2(\vartheta)-k^2}+\sum_{j=1}^{\infty}\sum_{s=1}^{\nu_*-\nu}\frac{Q_{s,j}^0(\vartheta)}{(\pi j)^2-k^2}\bigg)d\vartheta, \tag{4.21}$$

where

$$\begin{aligned} Q_{n,j}(\vartheta)&=\int_{\Gamma_*}\sum_{m\in\mathbb{Z}^d}Q(x+m)|\Psi_\alpha(\vartheta,x)|^2dx,\\ Q_{s,j}^0(\vartheta)&=\int_{\Gamma_*}\sum_{m\in\mathbb{Z}^d}Q(x+m)|\Psi_{s,j}^0(\vartheta,x)|^2dx. \end{aligned} \tag{4.22}$$

Proof. Due to Theorem 1.3 and (4.3) we have

$$\begin{aligned} &((\Delta_{MV}-k^2)^{-1}f,g)\\ &=\int_{\Gamma_*}dx\int_{\Gamma_*}dx'\int_{\mathbb{T}^d}\frac{d\vartheta}{(2\pi)^d}\sum_{m,\,m'\in\mathbb{Z}^d}e^{-i\langle m-m',\vartheta\rangle}\mathcal{R}_k(x,x',\vartheta)f(x+m)\overline{g}(x'+m'), \end{aligned}$$

where $f,g\in L^2(\Gamma)$ are compactly supported functions. Then this yields

$$\begin{aligned} &\operatorname{Tr}\big(|Q|^{\frac{1}{2}}(\Delta_{MV}-k^2)^{-1}Q^{\frac{1}{2}}\big)=\int_{\Gamma_*}dx\int_{\mathbb{T}^d}\frac{d\vartheta}{(2\pi)^d}\sum_{m\in\mathbb{Z}^d}\mathcal{R}_k(x,x,\vartheta)Q(x+m)\\ &=\sum_{j=0}^{\infty}\sum_{n=1}^{\nu}\int_{\Gamma_*}dx\int_{\mathbb{T}^d}\frac{d\vartheta}{(2\pi)^d}\sum_{m\in\mathbb{Z}^d}\frac{Q(x+m)|\Psi_\alpha(\vartheta,x)|^2}{z_{n,j}^2(\vartheta)-k^2}=\frac{1}{(2\pi)^d}\int_{\mathbb{T}^d}\sum_{j=0}^{\infty}\sum_{n=1}^{\nu}\frac{Q_\alpha(\vartheta)}{z_{n,j}^2(\vartheta)-k^2}d\vartheta. \end{aligned}$$

Similar arguments yield

$$\begin{aligned} &\operatorname{Tr}\big(|Q|^{\frac{1}{2}}(\Delta_{MD}-k^2)^{-1}Q^{\frac{1}{2}}\big)\\ &=\sum_{j=0}^{\infty}\sum_{s=1}^{\nu_*-\nu}\int_{\Gamma_*}dx\int_{\mathbb{T}^d}\frac{d\vartheta}{(2\pi)^d}\sum_{m\in\mathbb{Z}^d}\frac{Q(x+m)|\Psi_{s,j}^0(\vartheta,x)|^2}{(\pi j)^2-k^2}=\frac{1}{(2\pi)^d}\int_{\mathbb{T}^d}\sum_{j=0}^{\infty}\sum_{s=1}^{\nu_*-\nu}\frac{Q_{s,j}^0(\vartheta)}{(\pi j)^2-k^2}d\vartheta. \end{aligned}$$

Combining these results and using the identity

$$\operatorname{Tr}Y_0(k)=\operatorname{Tr}\big(|Q|^{\frac{1}{2}}(\Delta_{MV}-k^2)^{-1}Q^{\frac{1}{2}}\big)+\operatorname{Tr}\big(|Q|^{\frac{1}{2}}(\Delta_{MD}-k^2)^{-1}Q^{\frac{1}{2}}\big),$$

we obtain (4.21). ■

4.2. **Fredholm determinants.** In order to discuss Fredholm determinants we recall well-known facts [S05]. Let $\mathcal{H}$ be a Hilbert space endowed with an inner product (,) and a norm $\|\cdot\|$. Let $\mathcal{B}$ denote the class of bounded operators. Let $\mathbf{B}_1$ be the set of all trace class operators on $\mathcal{H}$ equipped with the trace norm $\|\cdot\|_{\mathbf{B}_1}$.

- Let $A,B\in\mathcal{B}$ and $AB,BA\in\mathbf{B}_1$. Then

$$\operatorname{Tr}AB=\operatorname{Tr}BA, \tag{4.23}$$

$$\det(I+AB)=\det(I+BA). \tag{4.24}$$

- Let $A,B\in\mathbf{B}_1$. Then

$$|\det(I+A)|\leqslant e^{\|A\|_{\mathbf{B}_1}}. \tag{4.25}$$

$$|\det(I+A)-\det(I+B)|\leqslant \|A-B\|_{\mathbf{B}_1}e^{1+\|A\|_{\mathbf{B}_1}+\|B\|_{\mathbf{B}_1}}. \tag{4.26}$$

Moreover, $I+A$ is invertible if and only if $\det(I+A)\neq 0$.

- Suppose for a domain $\mathscr{D}\subset\mathbb{C}$, the function $\Omega(\cdot)-I:\Omega\to\mathbf{B}_1$ is analytic and is invertible for any $z\in\mathscr{D}$. Then $F(z)=\det\Omega(z)$ is analytic and satisfies

$$F'(z)=F(z)\operatorname{Tr}\Omega^{-1}(z)\Omega'(z). \tag{4.27}$$

By Theorem 4.1, each operator $Y_0(k)\in\mathbf{B}_1$, $k\in\mathbb{C}_+$, and we can define the determinant

$$D(k)=\det(I+Y_0(k)),\qquad k\in\mathbb{C}_+. \tag{4.28}$$

Lemma 4.3. *Let $Q\in L^1(\Gamma)$. Then the determinant $D(k)$, $k\in\mathbb{C}_+$, is analytic in $k\in\mathbb{C}_+$. If, in addition, the constant $\mathcal{C}(Q,k)<1$, defined by (4.5), then*

$$\log D(k)=-\sum_{n=1}^{\infty}\frac{1}{n}\operatorname{Tr}\big(-Y_0(k)\big)^n, \tag{4.29}$$

where the series converges absolutely and uniformly, and for any $N\geqslant 0$ we have

$$\Big|\log D(k)+\sum_{n=1}^{N}\frac{\operatorname{Tr}(-Y_0(k))^n}{n}\Big|\leqslant\frac{\|Y_0(k)\|_{\mathbf{B}_1}^{N+1}}{(N+1)(1-\mathcal{C}(Q,k))}. \tag{4.30}$$

Proof. By Theorem 4.1, the $\mathbf{B}_1$-valued function $Y_0(k)$ is analytic in $\mathbb{C}_+$. The series (4.29) is standard, see [S05]. Under the condition $\|Y_0(k)\|_{\mathbf{B}_1}\leqslant\mathcal{C}(Q,k)$ the identity (4.29) gives (4.30). ■

Proof of Theorem 1.6. Due to (4.4), (4.5) we have (1.34). Then it is well known that there exist the wave operators $W_\pm$ and they are complete, see [RS78]. Moreover, S-operator given by (1.36) is unitary on $\mathscr{H}_{ac}(H_0)$ and the corresponding S-matrix $S(k)$ for almost all $k^2\in\sigma_{ac}(H_0)$ satisfies (1.37) and (1.38). ■

5. Lattice graph

We consider the lattice graph $\mathbb{L}^d=(V,\mathcal{E})$, where the vertex set and the edge set are given by

$$V=\mathbb{Z}^d,\qquad \mathcal{E}=\big\{(m,m+a_1),\ldots,(m,m+a_d),\quad\forall\, m\in\mathbb{Z}^d\big\}, \tag{5.1}$$

and $a_1,\ldots,a_d$ is the standard orthonormal basis, see Fig.2*a*. The "minimal" fundamental graph $\mathbb{L}^d_*=(V_*,\mathcal{E}_*)$ of the lattice $\mathbb{L}^d$ consists of one vertex v and d edge-loops $\mathbf{e}_1=\ldots=\mathbf{e}_d=(v,v)$ with indices $a_1,\ldots,a_d$, see Fig.2*b*. For each $\vartheta=(\vartheta_1,\ldots,\vartheta_d)\in\mathbb{T}^d$ the Floquet matrix $\Delta(\vartheta)$ degenerates to the scalar function

$$\Delta(\vartheta)=-\tfrac{1}{d}\,(\cos\vartheta_1+\ldots+\cos\vartheta_d), \tag{5.2}$$

which yields that the unique eigenvalue of $\Delta(\vartheta)$ and the corresponding normalized eigenfunction are given by

$$\lambda(\vartheta)=\Delta(\vartheta),\qquad \psi(\vartheta,v)=1. \tag{5.3}$$

The fiber operator $\Delta_M(\vartheta)$ acts on functions $y=(y_{\mathbf{e}})_{\mathbf{e}\in\mathcal{E}_*}\in L^2(\mathbb{L}^d_*)$ by $(\Delta_M(\vartheta)y)_{\mathbf{e}}=-y''_{\mathbf{e}}$, where $(y''_{\mathbf{e}})_{\mathbf{e}\in\mathcal{E}_*}\in L^2(\mathbb{L}^d_*)$ and y satisfies the quasi-periodic conditions:

$$y_{\mathbf{e}_1}(0)=\ldots=y_{\mathbf{e}_d}(0)=e^{-i\vartheta_1}y_{\mathbf{e}_1}(1)=\ldots=e^{-i\vartheta_d}y_{\mathbf{e}_d}(1), \tag{5.4}$$

$$y'_{\mathbf{e}_1}(0)+\ldots+y'_{\mathbf{e}_d}(0)-e^{-i\vartheta_1}y'_{\mathbf{e}_1}(1)-\ldots-e^{-i\vartheta_d}y'_{\mathbf{e}_d}(1)=0. \tag{5.5}$$

Indeed, substituting the indices $a_1,a_2,\ldots,a_d$ of the fundamental graph edges into the formulas (1.13), (1.14), we obtain the conditions (5.4) – (5.5).



FIGURE 2. $a)$ Lattice $\mathbb{L}^3$; $b)$ the fundamental graph $\mathbb{L}^3_*$.

Proposition 5.1. *Let $j\in\mathbb{N}$. Then for each $\vartheta\in\mathbb{T}^d\setminus\{0\}$ the operator $\Delta_M(\vartheta)$ has the eigenvalue $E^0_j=(\pi j)^2$ of multiplicity $d-1$ and the corresponding normalized eigenfunctions $\Psi^0_{s,j}=(\Psi^0_{s,j,\mathbf{e}})_{\mathbf{e}\in\mathcal{E}_*}$, $s=1,\ldots,d-1$, have the form*

$$\begin{aligned}&\Psi^0_{s,j,\mathbf{e}_p}(\vartheta,t)=\frac{\sqrt{2}\,J_p\sin(\pi jt)}{\sqrt{1+|C_{s,j}(\vartheta)|^2}}, \qquad \begin{cases}J_d=1, & J_s=-C_{s,j}(\vartheta)\\ J_p=0, & \forall\,p\neq s,d\end{cases},\\ &\textit{where}\quad C_{s,j}(\vartheta)=\frac{1-(-1)^je^{-i\vartheta_d}}{1-(-1)^je^{-i\vartheta_s}}.\end{aligned} \tag{5.6}$$

Proof. Since the number of the fundamental graph vertices $\nu=1$ and the number of the fundamental graph edges $\nu_*=d$, due to Theorem 1.2, for all $\vartheta\in\mathbb{T}^d\setminus\{0\}$ the eigenvalue $(\pi j)^2$ has multiplicity $\nu_*-\nu=d-1$ and all normalized eigenfunctions corresponding to the eigenvalue $(\pi j)^2$, have the form

$$\Psi^0_{s,j}=\big(\Psi^0_{s,j,\mathbf{e}}\big)_{\mathbf{e}\in\mathcal{E}_*},\qquad \Psi^0_{s,j,\mathbf{e}}(\vartheta,t)=X_{s,j,\mathbf{e}}(\vartheta)\sqrt{2}\,\sin(\pi jt)\,,\quad s=1,\ldots,d-1, \tag{5.7}$$

where $\big(X_{s,j,\mathbf{e}}(\vartheta)\big)_{\mathbf{e}\in\mathcal{E}_*}$ is a normalized solution of the equation

$$\sum_{p=1}^d x_p\big(1-(-1)^je^{-i\vartheta_p}\big)=0.$$

This equation has $d-1$ linearly independent solutions

$$\begin{aligned}
X_{1,j,\mathbf{e}_1}(\vartheta) = -\,C_{1,j}\,X_{1,j,\mathbf{e}_d}(\vartheta), \qquad & X_{1,j,\mathbf{e}_2}(\vartheta) = 0,\\
\ldots\,, \qquad X_{1,j,\mathbf{e}_{d-1}}(\vartheta) = 0, \qquad & X_{1,j,\mathbf{e}_d}(\vartheta);\\
X_{2,j,\mathbf{e}_1}(\vartheta) = 0, \qquad X_{2,j,\mathbf{e}_2}(\vartheta) = & -C_{2,j}\,X_{2,j,\mathbf{e}_d}(\vartheta),\\
\ldots\,, \qquad X_{2,j,\mathbf{e}_{d-1}}(\vartheta) = 0, \qquad & X_{2,j,\mathbf{e}_d}(\vartheta);\\
\ldots & \\
X_{d-1,j,\mathbf{e}_1}(\vartheta) = 0, \qquad X_{d-1,j,\mathbf{e}_2}(\vartheta) = 0, \qquad & \ldots\,,\\
X_{d-1,j,\mathbf{e}_{d-1}}(\vartheta) = -C_{d-1,j}X_{d-1,j,\mathbf{e}_d}(\vartheta), \qquad & X_{d-1,j,\mathbf{e}_d}(\vartheta),
\end{aligned} \tag{5.8}$$

where $C_{s,j} = C_{s,j}(\vartheta)$ are defined in (5.6). Each constant $X_{s,j,\mathbf{e}_d}(\vartheta)$, $s \in \mathbb{N}_{d-1}$, is determined by the condition $\|\Psi^0_{s,j}(\vartheta,\cdot)\|_{L^2(\mathbb{L}^d_*)} = 1$. Then (5.7), (5.8) give

$$\begin{aligned}
1 = \|\Psi^0_{s,j}(\vartheta,\cdot)\|^2_{L^2(\mathbb{L}^d_*)} &= \sum_{p=1}^{d}\int_0^1 \big|\Psi^0_{s,j,\mathbf{e}_p}(\vartheta,t)\big|^2 dt = \int_0^1 \big|\Psi^0_{s,j,\mathbf{e}_s}(\vartheta,t)\big|^2 dt + \int_0^1 \big|\Psi^0_{s,j,\mathbf{e}_d}(\vartheta,t)\big|^2 dt\\
&= 2\Big(\big|X_{s,j,\mathbf{e}_s}(\vartheta)\big|^2 + \big|X_{s,j,\mathbf{e}_d}(\vartheta)\big|^2\Big)\int_0^1 \sin^2(\pi j t)\,dt = \big(1+|C_{s,j}|^2\big)\big|X_{s,j,\mathbf{e}_d}(\vartheta)\big|^2,
\end{aligned}$$

which yields that the constant $X_{s,j,\mathbf{e}_d}(\vartheta)$ may be chosen in the form $X_{s,j,\mathbf{e}_d}(\vartheta) = \frac{1}{\sqrt{1+|C_{s,j}|^2}} > 0\,.$ Substituting this and (5.8) into (5.7), we obtain (5.6). ■

From (5.3) and Theorem 1.3 we deduce that the operator $\Delta_{MV}(\vartheta)$ on the metric graph $\mathbb{L}^d_*$, defined by (1.20), has the spectral representation

$$\Delta_{MV}(\vartheta) = \sum_{j=0}^{\infty} z_j^2(\vartheta)\mathcal{P}_j(\vartheta), \qquad \text{for all } \ \vartheta \in \mathbb{T}^d \setminus \{0\},$$

where its eigenvalues $z_j^2(\vartheta)$ and the corresponding normalized eigenfunctions $\Psi_j(\vartheta) = \big(\Psi_{j,\mathbf{e}}(\vartheta,t)\big)_{\mathbf{e}\in\mathcal{E}_*}$ are given by:

$$\begin{aligned}
&z_j(\vartheta) = \begin{cases} z(\vartheta) + \pi j, & j \text{ is even}\\ (\pi - z(\vartheta)) + \pi j, & j \text{ is odd}\end{cases}, \qquad z(\vartheta) = \arccos\big(-\lambda(\vartheta)\big) \in (0,\pi),\\
&\Psi_{j,\mathbf{e}_s}(\vartheta,t) = \tfrac{1}{\sqrt{d}\,\sin z(\vartheta)}\big(\sin(z_j(\vartheta)\,(1-t)) + e^{i\vartheta_s}\,\sin(z_j(\vartheta)\,t)\big), \qquad t \in [0,1].
\end{aligned} \tag{5.9}$$

Since $\lambda(\vartheta) = -\frac{1}{d}\,(\cos\vartheta_1 + \ldots + \cos\vartheta_d)$, we have

$$[\lambda^-,\lambda^+] = [-1,1], \qquad z^- = \arccos(-\lambda^-) = 0, \qquad z^+ = \arccos(-\lambda^+) = \pi.$$

Recall that $\mathbb{Z}_0 = \{0,1,2,\ldots\}$. Then, due to the formulas (1.26), the Laplacian Δ_M on $L^2(\mathbb{L}^d)$ has the spectrum given by

$$\begin{aligned}
\sigma(\Delta_M) = \sigma_{ac}(\Delta_M) \cup \sigma_{fb}(\Delta_M), \quad & \sigma_{fb}(\Delta_M) = \sigma_{fb}(\Delta_{MD}) = \sigma_D,\\
\sigma_{ac}(\Delta_M) = \bigcup_{j\in\mathbb{Z}_0} \sigma_j(\Delta_M) = [0,+\infty), \qquad & \sigma_j(\Delta_M) = \big[(\pi j)^2, (\pi+\pi j)^2\big].
\end{aligned} \tag{5.10}$$

6. Graphene lattice

We consider the hexagonal lattice $\mathbf{G}$ shown in Fig.3a. The periods a_1, a_2 of the lattice $\mathbf{G}$ and the fundamental vertex set $V_0 = \{v_1, v_2\}$ are also shown in the figure. The fundamental graph $\mathbf{G}_* = (V_*, \mathcal{E}_*)$, where $V_* = \{v_1, v_2\}$ consists of two vertices and three *multiple* edges $\mathbf{e}_1 = \mathbf{e}_2 = \mathbf{e}_3 = (v_1, v_2)$ (Fig.3b) with the indices $\tau(\mathbf{e}_1) = (1,0)$, $\tau(\mathbf{e}_2) = (0,1)$, $\tau(\mathbf{e}_3) = (0,0)$.

The Floquet matrix $\Delta(\vartheta)$, $\vartheta = (\vartheta_1, \vartheta_2) \in \mathbb{T}^2$, has the form

$$\Delta(\vartheta) = \begin{pmatrix} 0 & \phi(\vartheta) \\ \bar{\phi}(\vartheta) & 0 \end{pmatrix}, \qquad \phi(\vartheta) = -\tfrac{1}{3}\,(1 + e^{i\vartheta_1} + e^{i\vartheta_2}). \tag{6.1}$$

Then the eigenvalues of $\Delta(\vartheta)$ are given by

$$\lambda_1(\vartheta) = -\lambda_2(\vartheta) = -|\phi(\vartheta)|. \tag{6.2}$$



Figure 3. a) Graphene $\mathbf{G}$; b) the fundamental graph $\mathbf{G}_*$ of the graphene.

Proposition 6.1. *i) The fiber operator $\Delta_M(\vartheta)$ acts on $y = (y_\mathbf{e})_{\mathbf{e}\in\mathcal{E}_*} \in L^2(\mathbf{G}_*)$ by $(\Delta_M(\vartheta)y)_\mathbf{e} = -y''_\mathbf{e}$, where $(y''_\mathbf{e})_{\mathbf{e}\in\mathcal{E}_*} \in L^2(\mathbf{G}_*)$ and y satisfies the quasi-periodic conditions:*

$$y_{\mathbf{e}_1}(0) = y_{\mathbf{e}_2}(0) = y_{\mathbf{e}_3}(0), \qquad e^{-i\vartheta_1} y_{\mathbf{e}_1}(1) = e^{-i\vartheta_2} y_{\mathbf{e}_2}(1) = y_{\mathbf{e}_3}(1), \tag{6.3}$$

$$\begin{aligned} &y'_{\mathbf{e}_1}(0) + y'_{\mathbf{e}_2}(0) + y'_{\mathbf{e}_3}(0) = 0, \\ &e^{-i\vartheta_1} y'_{\mathbf{e}_1}(1) + e^{-i\vartheta_2} y'_{\mathbf{e}_2}(1) + y'_{\mathbf{e}_3}(1) = 0. \end{aligned} \tag{6.4}$$

ii) Let $j \in \mathbb{N}$. Then for each $\vartheta \in \mathbb{T}^2 \setminus \{0\}$ the operator $\Delta_M(\vartheta)$ has the simple eigenvalue $E_j^0 = (\pi j)^2$ and the corresponding normalized eigenfunction Ψ_j^0 has the form

$$\Psi_j^0 = \big(\Psi_{j,\mathbf{e}}^0\big)_{\mathbf{e}\in\mathcal{E}_*}, \qquad \Psi_{j,\mathbf{e}}^0(\vartheta, t) = X_\mathbf{e}(\vartheta)\sqrt{2}\,\sin(\pi j t)\,, \quad t \in [0,1], \tag{6.5}$$

where

$$X_{\mathbf{e}_1}(\vartheta) = -X_{\mathbf{e}_3}(\vartheta)\frac{1 - e^{-i\vartheta_2}}{e^{-i\vartheta_1} - e^{-i\vartheta_2}}\,, \qquad X_{\mathbf{e}_2}(\vartheta) = X_{\mathbf{e}_3}(\vartheta)\frac{1 - e^{-i\vartheta_1}}{e^{-i\vartheta_1} - e^{-i\vartheta_2}}\,, \tag{6.6}$$

$$X_{\mathbf{e}_3}(\vartheta) = \left(\frac{1 - \cos(\vartheta_1 - \vartheta_2)}{3 - \cos\vartheta_1 - \cos\vartheta_2 - \cos(\vartheta_1 - \vartheta_2)}\right)^{1/2}. \tag{6.7}$$

Proof. i) Substituting the indices $\tau(\mathbf{e}_1) = (1,0)$, $\tau(\mathbf{e}_2) = (0,1)$, $\tau(\mathbf{e}_3) = (0,0)$ of the fundamental graph edges into the formulas (1.13), (1.14), we obtain the conditions (6.3) – (6.4).

ii) Since the number of the fundamental graph vertices $\nu = 2$ and the number of the fundamental graph edges $\nu_* = 3$, due to Theorem 1.2, for all $\vartheta \in \mathbb{T}^2 \setminus \{0\}$ the eigenvalue $(\pi j)^2$ has multiplicity $\nu_* - \nu = 3 - 2 = 1$ and the unique normalized eigenfunction, corresponding to the eigenvalue $(\pi j)^2$, has the form (6.5), where $\big(X_{\mathbf{e}}(\vartheta)\big)_{\mathbf{e}\in\mathcal{E}_*}$ is the normalized solution of the system of two equations

$$\begin{aligned} x_{\mathbf{e}_1}+x_{\mathbf{e}_2}+x_{\mathbf{e}_3}&=0,\\ e^{-i\vartheta_1}x_{\mathbf{e}_1}+e^{-i\vartheta_2}x_{\mathbf{e}_2}+x_{\mathbf{e}_3}&=0, \end{aligned} \tag{6.8}$$

which yields (6.6). The constant $X_{\mathbf{e}_3}(\vartheta)$ is found using the condition $\|\Psi_j^0(\vartheta,\cdot)\|_{L^2(\mathbf{G}_*)}=1$. Using (6.5), (6.6), we have

$$1=\|\Psi_j^0(\vartheta,\cdot)\|^2_{L^2(\mathbf{G}_*)}=\sum_{s=1}^{3}\int_0^1\big|\Psi_{j,\mathbf{e}_s}(\vartheta,t)\big|^2dt=2\Big(\sum_{s=1}^{3}\big|X_{\mathbf{e}_s}(\vartheta)\big|^2\Big)\int_0^1\sin^2(\pi jt)\,dt$$
$$=\Big(1+\Big|\frac{1-e^{-i\vartheta_1}}{e^{-i\vartheta_1}-e^{-i\vartheta_2}}\Big|^2+\Big|\frac{1-e^{-i\vartheta_2}}{e^{-i\vartheta_1}-e^{-i\vartheta_2}}\Big|^2\Big)\big|X_{\mathbf{e}_3}(\vartheta)\big|^2=\Big(1+\frac{2-\cos\vartheta_1-\cos\vartheta_2}{1-\cos(\vartheta_1-\vartheta_2)}\Big)\big|X_{\mathbf{e}_3}(\vartheta)\big|^2,$$

which yields that the constant $X_{\mathbf{e}_3}(\vartheta)$ may be chosen in the form (6.7). ■

Proposition 6.2. *The operator $\Delta_{MV}(\vartheta)$ on $\mathbf{G}_*$, defined by (1.20), has the form*

$$\Delta_{MV}(\vartheta)=\sum_{j=0}^{\infty}z_{1,j}^2(\vartheta)\mathcal{P}_{1,j}(\vartheta)+\sum_{j=0}^{\infty}z_{2,j}^2(\vartheta)\mathcal{P}_{2,j}(\vartheta),\quad \textit{for all}\quad \vartheta\in\mathbb{T}^2\setminus\{0\}, \tag{6.9}$$

where its eigenvalues $z_{n,j}^2(\vartheta)$ and the corresponding normalized eigenfunctions $\Psi_{n,j}(\vartheta)=\big(\Psi_{n,j,\mathbf{e}}(\vartheta,t)\big)_{\mathbf{e}\in\mathcal{E}_*}$ have the form:

$$z_{n,j}(\vartheta)=\begin{cases} z_n(\vartheta)+\pi j, & j \text{ is even}\\ (\pi-z_n(\vartheta))+\pi j, & j \text{ is odd}\end{cases},\qquad z_n(\vartheta)=\arccos\big((-1)^{n+1}|\phi(\vartheta)|\big)\in(0,\pi), \tag{6.10}$$

$$\Psi_{n,j,\mathbf{e}_s}(\vartheta,t)=\tfrac{1}{\sqrt{3}\,\sin z_n(\vartheta)}\Big(\tfrac{(-1)^n\phi(\vartheta)}{|\phi(\vartheta)|}\sin(z_{n,j}(\vartheta)\,(1-t))+e^{i\langle\tau(\mathbf{e}_s),\vartheta\rangle}\sin(z_{n,j}(\vartheta)\,t)\Big), \tag{6.11}$$

$\phi(\vartheta)$ is defined in (6.1).

Proof. Using (1.21), (1.22), we obtain (6.9), (6.10). From (6.1) it follows that the normalized eigenfunction $\psi_n(\vartheta,\cdot)$, $n=1,2$, corresponding to the eigenvalue $\lambda_n(\vartheta)$ satisfies

$$\begin{aligned} -\lambda_n(\vartheta)\psi_n(\vartheta,v_1)+\phi(\vartheta)\psi_n(\vartheta,v_2)&=0,\\ |\psi_n(\vartheta,v_1)|^2+|\psi_n(\vartheta,v_2)|^2&=1, \end{aligned} \tag{6.12}$$

which yields

$$\psi_n(\vartheta,v_1)=\frac{\phi(\vartheta)}{\lambda_n(\vartheta)}\psi_n(\vartheta,v_2)=\frac{\phi(\vartheta)}{(-1)^n|\phi(\vartheta)|}\psi_n(\vartheta,v_2) \tag{6.13}$$

and we may choose $\psi_n(\vartheta,v_2)=\frac{1}{\sqrt{2}}$. Substituting this and (6.13) into (1.23) of Theorem 1.3, we obtain (6.11). ■

Corollary 6.3. *The Laplacian Δ_M on $L^2(\mathbf{G})$ has the spectrum given by*

$$\begin{aligned} \sigma(\Delta_M)&=\sigma_{ac}(\Delta_M)\cup\sigma_{fb}(\Delta_M),\\ \sigma_{ac}(\Delta_M)=[0,+\infty),\qquad \sigma_{fb}(\Delta_M)&=\sigma_{fb}(\Delta_{MD})=\sigma_D. \end{aligned} \tag{6.14}$$

Proof. The eigenvalues of $\Delta(\vartheta)$ is given by $\lambda_n(\vartheta) = \frac{(-1)^n}{3}|1 + e^{i\vartheta_1} + e^{i\vartheta_2}|$, $n = 1, 2$, which yields

$$[\lambda_1^-, \lambda_1^+] = \lambda_1(\mathbb{T}^2) = [-1, 0], \qquad [\lambda_2^-, \lambda_2^+] = \lambda_2(\mathbb{T}^2) = [0, 1]$$

and

$$z_1^- = \arccos(-\lambda_1^-) = 0, \qquad z_1^+ = \arccos(-\lambda_1^+) = \tfrac{\pi}{2},$$
$$z_2^- = \arccos(-\lambda_2^-) = \tfrac{\pi}{2}, \qquad z_2^+ = \arccos(-\lambda_2^+) = \pi.$$

Then, using the formulas (1.26), we obtain

$$\sigma_{1,j}(\Delta_M) = [E_{1,j}^-, E_{1,j}^+] = \begin{cases} \big[(\pi j)^2, \big(\frac{\pi}{2} + \pi j\big)^2\big], & j \text{ is even} \\ \big[\big(\frac{\pi}{2} + \pi j\big)^2, (\pi + \pi j)^2\big], & j \text{ is odd} \end{cases},$$

$$\sigma_{2,j}(\Delta_M) = [E_{2,j}^-, E_{2,j}^+] = \begin{cases} \big[\big(\frac{\pi}{2} + \pi j\big)^2, \big(\pi + \pi j\big)^2\big], & j \text{ is even} \\ \big[(\pi j)^2, \big(\frac{\pi}{2} + \pi j\big)^2\big], & j \text{ is odd} \end{cases}, \qquad j \in \mathbb{Z}_0 = \{0, 1, 2, \ldots\},$$

$$\sigma_{ac}(\Delta_M) = \bigcup_{n=1}^{2} \bigcup_{j \in \mathbb{Z}_0} \sigma_{n,j}(\Delta_M) = [0, +\infty), \quad \sigma_{fb}(\Delta_{MV}) = \varnothing, \quad \sigma_{fb}(\Delta_M) = \sigma_{fb}(\Delta_{MD}) = \sigma_D.$$

Thus, (6.14) has been proved. ■

7. Stanene lattice

Stanene is a topological insulator, theoretically predicted by Prof. Shoucheng Zhang's group at Stanford, which may display dissipationless current at its edges above room temperature [Z13]. It is composed of tin atoms arranged in a single layer, in a manner similar to graphene. The addition of fluorine atoms to the tin lattice could extend the critical temperature up to 100°C. This would make it practical for use in integrated circuits to make smaller, faster and more energy efficient computers. Stanene has a band gap, it is a semiconducting material. That makes it useful as material for use in a transistor, which must have a component that turns on and off. For more details see [Z13] and references therein.



Figure 4. a) Stanene $\mathbf{S}$; b) the fundamental graph $\mathbf{S}_*$ of the stanene.

A vertex of degree one is called an end vertex. An edge incident to an end vertex is called a pendant edge. The stanene lattice $\mathbf{S}$ is obtained from the hexagonal lattice $\mathbf{G}$ by adding a pendant edge at each vertex of $\mathbf{G}$ (see Fig.4a). We choose the orientations of the edges such as it is shown in Fig.4b. We consider the discrete and metric Laplacians on the stanene lattice

S. The fundamental graph $\mathbf{S}_*$ consists of 4 vertices v_1, v_2, v_3, v_4 with degrees $\varkappa_1 = \varkappa_2 = 4$, $\varkappa_3 = \varkappa_4 = 1$ and 5 oriented edges

$$\mathbf{e}_1 = \mathbf{e}_2 = \mathbf{e}_3 = (v_1, v_2), \qquad \mathbf{e}_4 = (v_1, v_3), \qquad \mathbf{e}_5 = (v_2, v_4).$$

The indices of the fundamental graph edges with respect to the fundamental vertex set $V_0 = \{v_1, v_2, v_3, v_4\}$, see Fig.4$a$, are given by

$$\tau(\mathbf{e}_1) = (1, 0), \qquad \tau(\mathbf{e}_2) = (0, 1), \qquad \tau(\mathbf{e}_3) = \tau(\mathbf{e}_4) = \tau(\mathbf{e}_5) = (0, 0). \tag{7.1}$$

For each $\vartheta = (\vartheta_1, \vartheta_2) \in \mathbb{T}^2$ the matrix $\Delta(\vartheta)$ has the form

$$\Delta(\vartheta) = -\frac{1}{2}\begin{pmatrix} 0 & b(\vartheta) & 1 & 0 \\ \overline{b}(\vartheta) & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}, \qquad b(\vartheta) = \frac{1}{2}\big(1 + e^{i\vartheta_1} + e^{i\vartheta_2}\big). \tag{7.2}$$

A direct calculation gives

$$\det\big(\Delta(\vartheta) - \lambda 1\!\!1_4\big) = \lambda^4 - \big(\tfrac{1}{2} + \tfrac{|b(\vartheta)|^2}{4}\big)\lambda^2 + \tfrac{1}{16}\,, \tag{7.3}$$

where $1\!\!1_4$ is the identity 4×4 matrix. The eigenvalues of the matrix $\Delta(\vartheta)$ are given by

$$\lambda_{1,2,3,4}(\vartheta) = \pm\frac{|b(\vartheta)|}{4} \pm \frac{\sqrt{|b(\vartheta)|^2 + 4}}{4}\,. \tag{7.4}$$

Proposition 7.1. *The spectrum of the discrete Laplacian* Δ *on the stanene lattice* $\mathbf{S}$ *has the form*

$$\begin{gathered} \sigma(\Delta) = \sigma_{ac}(\Delta) = \bigcup_{n=1}^{4} \sigma_n(\Delta) = \big[-1; -\tfrac{1}{4}\big] \cup \big[\tfrac{1}{4}\,; 1\big], \\ \sigma_1(\Delta) = [-1, -\tfrac{1}{2}], \qquad \sigma_2(\Delta) = [-\tfrac{1}{2}, -\tfrac{1}{4}], \qquad \sigma_3(\Delta) = [\tfrac{1}{4}, \tfrac{1}{2}], \qquad \sigma_4(\Delta) = [\tfrac{1}{2}, 1]. \end{gathered} \tag{7.5}$$

Proof. We need the following statement (see Theorem 4.3.8 in [HJ85]).

Let $B = \begin{pmatrix} A & y \\ y^* & a \end{pmatrix}$ *be a self-adjoint* $(\nu+1) \times (\nu+1)$ *matrix for some self-adjoint* $\nu \times \nu$ *matrix* A*, some real number* a *and some vector* $y \in \mathbb{C}^\nu$*. Denote by* $\lambda_1(A) \leqslant \ldots \leqslant \lambda_\nu(A)$ *the eigenvalues of* A*, arranged in increasing order, counting multiplicities. Then*

$$\lambda_1(B) \leqslant \lambda_1(A) \leqslant \lambda_2(B) \leqslant \ldots \leqslant \lambda_\nu(B) \leqslant \lambda_\nu(A) \leqslant \lambda_{\nu+1}(B).$$

Applying this statement to the matrix $\Delta(\vartheta)$, we obtain

$$\lambda_1(\vartheta) \leqslant -\tfrac{1}{2} \leqslant \lambda_2(\vartheta) \leqslant 0 \leqslant \lambda_3(\vartheta) \leqslant \tfrac{1}{2} \leqslant \lambda_4(\vartheta), \qquad \forall\, \vartheta \in \mathbb{T}^2. \tag{7.6}$$

Since

$$\lambda_1(0) = -1, \qquad \lambda_1\big(\tfrac{2\pi}{3}\,, -\tfrac{2\pi}{3}\big) = -\tfrac{1}{2}\,,$$

using (7.6) and the fact that the spectrum of a bipartite graph is symmetric with respect to the point 0, we have

$$\sigma_1 = \lambda_1(\mathbb{T}^2) = [-1, -\tfrac{1}{2}], \qquad \sigma_4 = \lambda_4(\mathbb{T}^2) = -\lambda_1(\mathbb{T}^2) = [\tfrac{1}{2}, 1]. \tag{7.7}$$

From (7.3) and (7.6) it follows that

$$\lambda_1(\vartheta)\lambda_2(\vartheta) = \tfrac{1}{4}\,. \tag{7.8}$$

This and (7.7) yield

$$\sigma_2 = \lambda_2(\mathbb{T}^2) = \big[-\tfrac12, -\tfrac14\big], \qquad \sigma_3 = \lambda_3(\mathbb{T}^2) = -\lambda_2(\mathbb{T}^2) = \big[\tfrac14, \tfrac12\big]. \tag{7.9}$$

Thus, (7.5) has been proved. ■

Proposition 7.2. *i) The fiber operator $\Delta_M(\vartheta)$ acts on $y = (y_{\mathbf{e}})_{\mathbf{e}\in\mathcal{E}_*} \in L^2(\mathbf{S}_*)$ by $(\Delta_M(\vartheta)y)_{\mathbf{e}} = -y''_{\mathbf{e}}$, where $(y''_{\mathbf{e}})_{\mathbf{e}\in\mathcal{E}_*} \in L^2(\mathbf{S}_*)$ and y satisfies the quasi-periodic conditions:*

$$\begin{aligned} y_{\mathbf{e}_1}(0) = y_{\mathbf{e}_2}(0) = y_{\mathbf{e}_3}(0) = y_{\mathbf{e}_4}(0),\\ e^{-i\vartheta_1}y_{\mathbf{e}_1}(1) = e^{-i\vartheta_2}y_{\mathbf{e}_2}(1) = y_{\mathbf{e}_3}(1) = y_{\mathbf{e}_5}(1), \end{aligned} \tag{7.10}$$

$$\begin{aligned} y'_{\mathbf{e}_1}(0) + y'_{\mathbf{e}_2}(0) + y'_{\mathbf{e}_3}(0) + y'_{\mathbf{e}_4}(0) = 0,\\ e^{-i\vartheta_1}y'_{\mathbf{e}_1}(1) + e^{-i\vartheta_2}y'_{\mathbf{e}_2}(1) + y'_{\mathbf{e}_3}(1) - y'_{\mathbf{e}_5}(1) = 0,\\ y'_{\mathbf{e}_4}(1) = 0, \qquad y'_{\mathbf{e}_5}(1) = 0. \end{aligned} \tag{7.11}$$

ii) Let $j \in \mathbb{N}$. Then for each $\vartheta \in \mathbb{T}^2\setminus\{0\}$ the operator $\Delta_M(\vartheta)$ has only the simple eigenvalue $E^0_j = (\pi j)^2$ and the corresponding normalized eigenfunction $\Psi^0_j = \big(\Psi^0_{j,\mathbf{e}}\big)_{\mathbf{e}\in\mathcal{E}_*}$ has the form

$$\Psi^0_{j,\mathbf{e}_s}(\vartheta,t) = X_{\mathbf{e}_s}(\vartheta)\sqrt{2}\sin(\pi j t)\,, \quad s = 1,2,3, \qquad \Psi^0_{j,\mathbf{e}_4}(\vartheta,t) = \Psi^0_{j,\mathbf{e}_5}(\vartheta,t) \equiv 0, \tag{7.12}$$

where the constants $X_{\mathbf{e}_s}(\vartheta)$, $s = 1,2,3$, are defined by (6.6), (6.7).

Proof. i) Substituting the indices (7.1) of the fundamental graph edges into the formulas (1.13), (1.14), we obtain the conditions (7.10) – (7.11).

ii) Since the number of the fundamental graph vertices $\nu = 4$ and the number of the fundamental graph edges $\nu_* = 5$, due to Theorem 1.2, for all $\vartheta \in \mathbb{T}^2\setminus\{0\}$ the eigenvalue $(\pi j)^2$ has multiplicity $\nu_* - \nu = 5 - 4 = 1$ and the unique normalized eigenfunction, corresponding to the eigenvalues $(\pi j)^2$, has the form

$$\Psi^0_{j,\mathbf{e}}(\vartheta,t) = X_{\mathbf{e}}(\vartheta)\sqrt{2}\sin(\pi j t)\,, \tag{7.13}$$

where $\big(X_{\mathbf{e}}(\vartheta)\big)_{\mathbf{e}\in\mathcal{E}_*}$ is the normalized solution of the system of 4 equations

$$\begin{aligned} x_{\mathbf{e}_1} + x_{\mathbf{e}_2} + x_{\mathbf{e}_3} + x_{\mathbf{e}_4} = 0,\\ e^{-i\vartheta_1}x_{\mathbf{e}_1} + e^{-i\vartheta_2}x_{\mathbf{e}_2} + x_{\mathbf{e}_3} - (-1)^j x_{\mathbf{e}_5} = 0, \qquad x_{\mathbf{e}_4} = 0, \qquad x_{\mathbf{e}_5} = 0, \end{aligned} \tag{7.14}$$

which yields the system (6.8) for $X_{\mathbf{e}_s}$, $s = 1,2,3$. The unique normalized solution of (6.8) is given by (6.6), (6.7). Thus, (7.12) has been proved. ■

Proposition 7.3. *The operator $\Delta_{MV}(\vartheta)$ on $\mathbf{S}_*$, defined by (1.20), has the form*

$$\Delta_{MV}(\vartheta) = \sum_{n=1}^{4}\sum_{j=0}^{\infty} z^2_{n,j}(\vartheta)\mathcal{P}_{n,j}(\vartheta), \quad \textit{for all} \quad \vartheta \in \mathbb{T}^2\setminus\{0\}, \tag{7.15}$$

where its eigenvalues $z^2_{n,j}(\vartheta)$ and the corresponding normalized eigenfunctions $\Psi_{n,j}(\vartheta) = \big(\Psi_{n,j,\mathbf{e}}(\vartheta,t)\big)_{\mathbf{e}\in\mathcal{E}_*}$ have the form:

$$z_{n,j}(\vartheta) = \begin{cases} z_n(\vartheta) + \pi j, & j \textit{ is even}\\ (\pi - z_n(\vartheta)) + \pi j, & j \textit{ is odd} \end{cases}, \qquad z_n(\vartheta) = \arccos\big(-\lambda_n(\vartheta)\big) \in (0,\pi), \tag{7.16}$$

$$\begin{aligned}
\Psi_{n,j,\mathbf{e}_s}(\cdot,t) &= \tfrac{1}{\sqrt{2}\ \sin z_n(\cdot)}\Big(\psi_n(\cdot,v_1)\sin(z_{n,j}(\cdot)\,(1-t))+\psi_n(\cdot,v_2)e^{i\langle\tau(\mathbf{e}_s),\,\cdot\rangle}\ \sin(z_{n,j}(\cdot)\,t)\Big),\\
&\qquad s=1,2,3,\\
\Psi_{n,j,\mathbf{e}_4}(\cdot,t) &= \tfrac{\sqrt{2}}{\sin z_n(\cdot)}\Big(\tfrac{\psi_n(\cdot,v_1)}{2}\ \sin(z_{n,j}(\cdot)\,(1-t))+\psi_n(\cdot,v_3)\sin(z_{n,j}(\cdot)\,t)\Big),\\
\Psi_{n,j,\mathbf{e}_5}(\cdot,t) &= \tfrac{\sqrt{2}}{\sin z_n(\cdot)}\Big(\tfrac{\psi_n(\cdot,v_2)}{2}\sin(z_{n,j}(\cdot)\,(1-t))+\psi_n(\cdot,v_4)\sin(z_{n,j}(\cdot)\,t)\Big),
\end{aligned} \tag{7.17}$$

$$\begin{aligned}
\psi_n(\cdot,v_1) &= \Big(\tfrac{2\lambda_n^2(\cdot)}{1+4\lambda_n^2(\cdot)}\Big)^{1/2}, \qquad \psi_n(\cdot,v_2)=-\tfrac{1}{b(\cdot)}\big(2\lambda_n(\cdot)-\tfrac{1}{2\lambda_n(\cdot)}\big)\psi_n(\cdot,v_1),\\
\psi_n(\cdot,v_3) &= -\tfrac{1}{2\lambda_n(\cdot)}\,\psi_n(\cdot,v_1), \qquad \psi_n(\cdot,v_4)=\tfrac{1}{2\lambda_n(\cdot)b(\cdot)}\big(2\lambda_n(\cdot)-\tfrac{1}{2\lambda_n(\cdot)}\big)\psi_n(\cdot,v_1).
\end{aligned} \tag{7.18}$$

Proof. Using (1.21), (1.22), (1.23), we obtain (7.15), (7.16), (7.17).

For each $\vartheta\in\mathbb{T}^2$ the Floquet matrix $\Delta(\vartheta)$ has the form (7.2) and the eigenvalues of $\Delta(\vartheta)$ are given by (7.4). Then the eigenfunction $\psi_n(\vartheta,\cdot)$, corresponding to the eigenvalue $\lambda_n(\vartheta)$ satisfies

$$\begin{aligned}
&\lambda_n(\cdot)\psi_n(\cdot,v_1)+\tfrac12\,b(\cdot)\,\psi_n(\cdot,v_2)+\tfrac12\,\psi_n(\cdot,v_3)=0, \qquad \lambda_n(\cdot)\psi_n(\cdot,v_3)+\tfrac12\,\psi_n(\cdot,v_1)=0,\\
&\lambda_n(\cdot)\psi_n(\cdot,v_2)+\tfrac12\,\bar b(\cdot)\,\psi_n(\cdot,v_1)+\tfrac12\,\psi_n(\cdot,v_4)=0, \qquad \lambda_n(\cdot)\psi_n(\cdot,v_4)+\tfrac12\,\psi_n(\cdot,v_2)=0,
\end{aligned} \tag{7.19}$$

which are normalized by $\sum\limits_{s=1}^{4}|\psi_n(\cdot,v_s)|^2=1$. Using (7.3), we obtain that

$$|b(\vartheta)|=\Big|2\lambda_n(\vartheta)-\tfrac{1}{2\lambda_n(\vartheta)}\Big|. \tag{7.20}$$

Solving the system of the equations (7.19) and applying (7.20), we have (7.18). ■

Corollary 7.4. *The Laplacian Δ_M on $L^2(\mathbf{S})$ has the spectrum given by*

$$\begin{aligned}
\sigma(\Delta_M) &= \sigma_{ac}(\Delta_M)\cup\sigma_{fb}(\Delta_M), \qquad \sigma_{fb}(\Delta_M)=\sigma_{fb}(\Delta_{MD})=\sigma_D,\\
\sigma_{ac}(\Delta_M) &= \big[0,r_+^2\big]\cup_{j\in\mathbb{Z}_0}\big[\big(r_-+2\pi j\big)^2,\big(-r_-+2\pi(j+1)\big)^2\big],\\
&\quad \cup_{j\in\mathbb{Z}_0}\big[\big(-r_++2\pi(j+1)\big)^2,\big(r_++2\pi(j+1)\big)^2\big],
\end{aligned} \tag{7.21}$$

where $r_\pm=\arccos\big(\pm\frac14\big)$.

Proof. Due to Proposition 7.1, we have that the bands $[\lambda_n^-,\lambda_n^+]=\lambda_n(\mathbb{T}^2)$ satisfy

$$[\lambda_1^-,\lambda_1^+]=[-1,-\tfrac12], \qquad [\lambda_2^-,\lambda_2^+]=[-\tfrac12,-\tfrac14], \qquad [\lambda_3^-,\lambda_3^+]=[\tfrac14,\tfrac12], \qquad [\lambda_4^-,\lambda_4^+]=[\tfrac12,1],$$

and

$$\begin{aligned}
z_1^- &= \arccos(-\lambda_1^-)=0, & z_1^+ &= \arccos(-\lambda_1^+)=\tfrac{\pi}{3},\\
z_2^- &= \arccos(-\lambda_2^-)=\tfrac{\pi}{3}, & z_2^+ &= \arccos(-\lambda_2^+)=\arccos\big(\tfrac14\big),\\
z_3^- &= \arccos(-\lambda_3^-)=\arccos\big(-\tfrac14\big), & z_3^+ &= \arccos(-\lambda_3^+)=\tfrac{2\pi}{3},\\
z_4^- &= \arccos(-\lambda_4^-)=\tfrac{2\pi}{3}, & z_4^+ &= \arccos(-\lambda_4^+)=\pi.
\end{aligned}$$

Then, using the formulas (1.26), we obtain

$$\sigma_{1,j}(\Delta_M)=[E_{1,j}^-,E_{1,j}^+]=\begin{cases}\big[(\pi j)^2,\big(\frac{\pi}{3}+\pi j\big)^2\big], & j\text{ is even}\\ \big[\big(\frac{2\pi}{3}+\pi j\big)^2,(\pi+\pi j)^2\big], & j\text{ is odd}\end{cases},$$

$$\sigma_{2,j}(\Delta_M)=[E_{2,j}^-,E_{2,j}^+]=\begin{cases}\big[\big(\frac{\pi}{3}+\pi j\big)^2,\big(\arccos\big(\frac{1}{4}\big)+\pi j\big)^2\big], & j\text{ is even}\\ \big[(\pi-\arccos\big(\frac{1}{4}\big)+\pi j)^2,\big(\frac{2\pi}{3}+\pi j\big)^2\big], & j\text{ is odd}\end{cases},$$

$$\sigma_{3,j}(\Delta_M)=[E_{3,j}^-,E_{3,j}^+]=\begin{cases}\big[\big(\arccos\big(-\frac{1}{4}\big)+\pi j\big)^2,\big(\frac{2\pi}{3}+\pi j\big)^2\big], & j\text{ is even}\\ \big[\big(\frac{\pi}{3}+\pi j\big)^2,(\pi-\arccos\big(-\frac{1}{4}\big)+\pi j)^2\big], & j\text{ is odd}\end{cases},$$

$$\sigma_{4,j}(\Delta_M)=[E_{4,j}^-,E_{4,j}^+]=\begin{cases}\big[(\frac{2\pi}{3}+\pi j)^2,\big(\pi+\pi j\big)^2\big], & j\text{ is even}\\ \big[\big(\pi j\big)^2,(\frac{\pi}{3}+\pi j)^2\big], & j\text{ is odd}\end{cases},$$

for all $j\in\mathbb{Z}_0=\{0,1,2,\dots\}$. Substituting these identities into the formula

$$\sigma_{ac}(\Delta_M)=\bigcup_{n=1}^{4}\bigcup_{j\in\mathbb{Z}_0}\sigma_{n,j}(\Delta_M),$$

we obtain the third identity in (7.21). Since $\sigma_{fb}(\Delta_{MV})=\varnothing$, we have the second formula in (7.21). ■

Acknowledgments. Our study was supported by the RSF grant No 15-11-30007. We would like to thank Olaf Post (Germany) for thorough reading of the manuscript and helpful remarks.

References

[A12] Ando, K. Inverse scattering theory for discrete Schrödinger operators on the hexagonal lattice, Ann. Henri Poincaré, 14 (2013), 347–383.

[B85] J. von Below, A characteristic equation associated to an eigenvalue problem on c^2-networks, Linear Algebra Appl., 71 (1985), 309-325.

[BK13] Berkolaiko, G.; Kuchment, P. Introduction to Quantum Graphs, Mathematical Surveys and Monographs, V. 186 AMS, 2013.

[BK62] Birman, M.; Krein, M. On the theory of wave operators and scattering operators, Dokl. Akad. Nauk SSSR, Ser. Mat., 144 (1962), 475-478; English transl.: Soviet Math. Dokl., 3 (1962), 740–744.

[BS99] Boutet de Monvel, A.; Sahbani, J. On the spectral properties of discrete Schrödinger operators : (The multi-dimensional case), Review in Math. Phys., 11 (1999), 1061–1078.

[BGP08] Brüning, J.; Geyler, V.; Pankrashkin, K. Spectra of self-adjoint extensions and applications to solvable Schrödinger operators, Rev. Math. Phys., 20 (2008), 1–70.

[C97] Cattaneo, C. The spectrum of the continuous Laplacian on a graph, Monatsh. Math. 124 (1997), 215–235.

[Ch97] Chung, F. Spectral graph theory, AMS, Providence, Rhode. Island, 1997.

[CDS95] Cvetkovic, D.; Doob, M.; Sachs, H. Spectra of graphs. Theory and applications. Third edition. Johann Ambrosius Barth, Heidelberg, 1995.

[CDGT88] Cvetkovic, D.; Doob, M.; Gutman, I.; Torgašev, A. Recent results in the theory of graph spectra. Annals of Discrete Mathematics, 36. North-Holland Publishing Co., Amsterdam, 1988.

[GKT93] Gieseker, D.; Knörrer, H.; Trubowitz, E. The geometry of algebraic Fermi curves. Perspectives in Mathematics, 14. Academic Press, Inc., Boston, MA, 1993.

[HJ85] Horn, R; Johnson, C. Matrix analysis. Cambridge University Press, 1985.

[HN09] Higuchi, Y.; Nomura, Y. Spectral structure of the Laplacian on a covering graph, European J. Combin., 30 (2009), no. 2, 570–585.

[HS04] Higuchi, Y.; Shirai, T. Some spectral and geometric properties for infinite graphs, AMS Contemp. Math., 347 (2004), 29–56.

[HS02] Hundertmark, D.; Simon, B. Lieb-Thirring Inequalities for Jacobi Matrices, Jour. Approx. Theory, 118 (2002), 106–130.

[IK12] Isozaki, H.; Korotyaev, E. Inverse problems, trace formulae for discrete Schrödinger operators, Ann. Henri Poincare, 13 (2012), no 4, 751–788.

[IM14] Isozaki, H.; Morioka, H. A Rellich type theorem for discrete Schrödinger operators, Inverse Probl. Imaging, 8 (2014), no. 2, 475–489.

[KK10] Korotyaev, E.; Kutsenko, A. Zigzag nanoribbons in external electric fields, Asympt. Anal., 66 (2010), no 3-4, 187–206.

[KS14a] Korotyaev, E.; Saburova, N., Schrödinger operators on periodic discrete graphs, J. Math. Anal. Appl., 420 (2014), No 1, 576–611.

[KS14b] Korotyaev, E.; Saburova, N. Spectral band bracketing for Laplacians on periodic metric graphs, preprint arXiv:1406.7523.

[KS15a] Korotyaev, E.; Saburova, N. Spectral band localization for Schrödinger operators on periodic graphs, Proc. Amer. Math. Soc., 143 (2015), 3951–3967.

[KS15b] Korotyaev, E.; Saburova, N. Estimates of bands for Laplacians on periodic equilateral metric graphs, to be published in Proc. Amer. Math. Soc.

[KS15c] Korotyaev, E.; Saburova, N. Effective masses for Laplacians on periodic graphs, preprint arXiv:1502.02989.

[LP08] Lledó, F.; Post, O. Eigenvalue bracketing for discrete and metric graphs, J. Math. Anal. Appl., 348 (2008), 806–833.

[Na14] Nakamura, S., Modified wave operators for discrete Schrödinger operators with long-range perturbations, preprint arXiv:1403.2795, 2014.

[N77] Newton, R. Noncentral potentials: The generalized Levinson theorem and the structure of the spectrum, J. Math. Phys., 18 (1977), 1348–1357.

[Pa12] Pankrashkin, K. Unitary dimension reduction for a class of self-adjoint extensions with applications to graph-like structures, J. Math. Anal. Appl., 396 (2012), no. 2, 640–655.

[Pa13] Pankrashkin, K. An example of unitary equivalence between self-adjoint extensions and their parameters, J. Funct. Anal. 265 (2013), 2910–2936.

[P12] Post, O. Spectral analysis on graph-like spaces. Lecture Notes in Mathematics, 2039. Springer, Heidelberg, 2012.

[RS78] Reed, M.; Simon, B. Methods of modern mathematical physics, Vol.IV, Analysis of operators, Academic Press, New York, 1978.

[RR07] Rabinovich, V. S.; Roch, S. Essential spectra of difference operators on $\mathbf{Z}^n$-periodic graphs, J. Phys. A: Math. Theor., 40 (2007), no. 33, 10109–10128.

[RoS09] Rosenblum, G.; Solomjak, M. On the spectral estimates for the Schrödinger operator on $\mathbb{Z}^d$, $d \geqslant 3$, Problems in Mathematical Analysis, No. 41, J. Math. Sci. N. Y. 159 (2009), No. 2, 241–263.

[SV01] Shaban, W.; Vainberg, B. Radiation conditions for the difference Schrödinger operators, J. Appl. Anal., 80 (34) (2001) 525-556.

[S05] Simon, B. Trace Ideals and Their Applications, Mathematical Surveys and Monographs, V. 120 AMS, 2005.

[Z13] Xu, Y.; Yan, B.; Zhang, H. J.; Wang, J.; Xu, G.; Tang, P.; Duan, W.; Zhang, S.C. Large-Gap Quantum Spin Hall Insulators in Tin Films, Phys. Review Letters, 111(2013), 136804.

Mathematical Physics Department, Faculty of Physics, Ulianovskaya 2, St. Petersburg State University, St. Petersburg, 198504, korotyaev@gmail.com,

Department of Mathematical Analysis, Algebra and Geometry, Institute of Mathematics, Information and Space Technologies, Uritskogo St. 68, Northern (Arctic) Federal University, Arkhangelsk, 163002, n.saburova@gmail.com